


Recombinant HA Tag Antibody [clone RM305] (R20304)

| Catalog No. | Formulation | Size |
|--------------|---|--------|
| R20304-100UG | 1 mg/ml in PBS with 50% glycerol, 1% BSA and 0.09% sodium azide | 100 ug |


Recombinant **RABBIT MONOCLONAL**

Bulk quote request

| | |
|--------------------|--|
| Availability | 1-3 business days |
| Species Reactivity | NA |
| Format | Purified |
| Host | Rabbit |
| Clonality | Recombinant Rabbit Monoclonal |
| Isotype | Rabbit IgG |
| Clone Name | RM305 |
| Purity | Protein A purified from animal origin-free supernatant |
| UniProt | N/A |
| Applications | Western Blot : 0.1 ug/ml-1 ug/ml Immunoprecipitation : 0.5 ug/ml-2 ug/ml Immunocytochemistry : 0.1 ug/ml-1 ug/ml Immunohistochemistry : 0.1 ug/ml-0.5 ug/ml |
| Limitations | This recombinant HA Tag antibody is available for research use only. |



Western blot testing of recombinant HA Tag antibody (0.1ug/ml) with lysate from HEK293 cells transfected (+) or untransfected (-) with an HA-tag KRas DNA construct.



IHC staining of formalin fixed and paraffin embedded human LNCaP cells overexpressing HA-tag Bag1 protein, using HA-Tag antibody at 0.01 ug/ml.

Description

This antibody reacts to recombinant proteins containing a HA-Tag fused to either the amino or carboxy terminus. No cross reactivity with other endogenous protein.

Application Notes

The stated application concentrations are suggested starting points. Titration of the recombinant HA Tag antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

A peptide corresponding to HA-tag (Human influenza hemagglutinin amino acids 98-106) was used as the immunogen for the recombinant HA Tag antibody.

Storage

Store the recombinant HA Tag antibody at -20oC.