


## Human IgG4 Fc Antibody for ELISA / Anti-IgG4 Fc Detection Antibody [clone RM217] (R20191)

Catalog No.	Formulation	Size
R20191-100UG	1 mg/ml in PBS with 50% glycerol, 1% BSA and 0.09% sodium azide	100 ug


Recombinant **RABBIT MONOCLONAL**

[Bulk quote request](#)


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Host</b>	Rabbit
<b>Clonality</b>	Recombinant Rabbit Monoclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Name</b>	RM217
<b>Purity</b>	Protein A purified from animal origin-free supernatant
<b>UniProt</b>	P01861
<b>Gene ID</b>	3503
<b>Applications</b>	ELISA : 50ng/well-200ng/well (Capture); 0.05-0.2ug/ml (Detection)
<b>Limitations</b>	This Human IgG4 Fc Antibody for ELISA / Anti-IgG4 Fc Detection Antibody is available for research use only.




Human IgG4 Fc Antibody for ELISA Sandwich ELISA Human Plasma. Sandwich ELISA analysis demonstrates that clone RM217 functions effectively as a capture antibody for IgG4 / IGHG4 in human plasma, with signal intensity decreasing proportionally with dilution, indicating concentration-dependent detection. Captured IgG4 was detected using a biotinylated anti-human light chains (kappa + lambda) antibody (clone RM129), followed by alkaline phosphatase-conjugated streptavidin for signal development. This assay supports selective Fc-region detection of IgG4 and highlights the utility of this Human IgG4 Fc Antibody for ELISA / Anti-IgG4 Fc Detection Antibody in subclass-specific ELISA workflows.



Human IgG4 Fc Antibody for ELISA Sandwich ELISA Human IgG4. Sandwich ELISA analysis using purified human IgG4 demonstrates that clone RM217 functions effectively as a capture antibody for IgG4 / IGHG4, with signal intensity decreasing proportionally with antigen concentration, indicating strong and concentration-dependent detection. Captured IgG4 was detected using a biotinylated anti-human light chains (kappa + lambda) antibody (clone RM129), followed by alkaline phosphatase-conjugated streptavidin for signal development. This assay supports selective Fc-region detection of IgG4 and highlights the utility of this Human IgG4 Fc Antibody for ELISA / Anti-IgG4 Fc Detection Antibody in subclass-specific ELISA applications.



Human IgG4 Fc Antibody for ELISA Subclass Specificity Analysis. ELISA analysis of human IgG subclasses demonstrates that clone RM217 selectively recognizes IgG4 / IGHG4, with strong signal observed for IgG4 and no detectable binding to IgG1, IgG2, or IgG3. This result confirms Fc-region-specific recognition and strict subclass selectivity. The Human IgG4 Fc Antibody for ELISA / Anti-IgG4 Fc Detection Antibody enables accurate IgG4 detection in mixed immunoglobulin samples and supports high-specificity ELISA workflows requiring precise subclass discrimination.



suited for use in direct, indirect, and sandwich ELISA formats requiring high specificity and sensitivity.

Detection of IgG4 is widely applied in immunology research, allergy studies, and therapeutic antibody development. IgG4 is often associated with regulatory immune responses and long-term antigen exposure, and its accurate measurement provides important insight into antibody subclass distribution, immune tolerance, and biologic drug responses. This antibody supports these applications by enabling sensitive and specific detection of the IgG4 Fc region in ELISA-based systems requiring precise subclass discrimination, low background, and consistent performance.

This antibody is part of a broader [immunoglobulin detection antibody collection](#), including reagents for Ig classes and light chains across multiple species and immunoassay formats.

## Application Notes

The stated application concentrations are suggested starting points. Titration of the Human IgG4 Fc Antibody for ELISA / Anti-IgG4 Fc Detection Antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

Peptide corresponding to the Fc region of hIgG4 was used as the immunogen for this recombinant Human IgG4 Fc antibody.

## Storage

Store the recombinant Human IgG4 Fc antibody at -20oC (with glycerol) or aliquot and store at -20oC (without glycerol).

## Alternate Names

Anti-IgG4 Fc antibody, IgG4 Fc detection antibody, Human IgG4 Fc antibody, IgG4 constant region antibody, Fc-specific IgG4 antibody