


Recombinant Interleukin 2 Antibody [clone IL2/7051R] (V9562)

| Catalog No. | Formulation | Size |
|----------------|---|--------|
| V9562-100UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 100 ug |
| V9562-20UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 20 ug |
| V9562SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free | 100 ug |


Recombinant **RABBIT MONOCLONAL**

[Bulk quote request](#)

| | |
|---------------------------|---|
| Availability | 1-3 business days |
| Species Reactivity | Human |
| Format | Purified |
| Clonality | Recombinant Rabbit Monoclonal |
| Isotype | Rabbit IgG |
| Clone Name | IL2/7051R |
| Purity | Protein A/G affinity |
| UniProt | P60568 |
| Localization | Secreted |
| Applications | Immunohistochemistry (FFPE) : 1-2ug/ml |
| Limitations | This recombinant Interleukin 2 antibody is available for research use only. |



IHC staining of FFPE human colon carcinoma tissue with recombinant Interleukin 2 antibody (clone IL2/7051R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free recombinant Interleukin 2 antibody (clone IL2/7051R) as confirmation of integrity and purity.

Description

Interleukin-2 (IL-2) is a T cell stimulatory cytokine that induces T cell proliferation and NK cell proliferation and activation.

Application Notes

Optimal dilution of the recombinant Interleukin 2 antibody should be determined by the researcher.

Immunogen

A portion of amino acids 21-153 was used as the immunogen for the recombinant Interleukin 2 antibody.

Storage

Aliquot the recombinant Interleukin 2 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.