


Recombinant ITGAE Antibody / Integrin alpha E / CD103 [clone rITGAE/2063] (V8433)

Catalog No.	Formulation	Size
V8433-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V8433-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V8433SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant MOUSE MONOCLONAL

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Mouse Monoclonal
Isotype	Mouse IgG1, kappa
Clone Name	rITGAE/2063
Purity	Protein G affinity chromatography
UniProt	P38570
Localization	Cell surface and cytoplasm
Applications	Immunohistochemistry (FFPE): 1-2ug/ml for 30 minutes at RT
Limitations	This recombinant ITGAE antibody is available for research use only.



IHC staining of FFPE human tonsil with recombinant ITGAE antibody (clone rITGAE/2063). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free recombinant ITGAE antibody (clone rITGAE/2063) as confirmation of integrity and purity.

Description

This MAb recognizes a protein of 150kDa, identified as CD103, which is the alpha-E integrin subunit of the heterodimeric alpha-E beta-7 (aEb7) integrin belonging to a small beta-7 integrin subfamily. CD103 is expressed on more than 95% of intraepithelial CD8+ cells and on 40% of mucosa-associated T cells, whereas less than 2% of resting blood lymphocytes are CD103-positive. In several malignant conditions, such as T-cell lymphomas and hairy cell leukemia (HCL), the cells express CD103. Antibody to CD103 is an extremely useful addition to the IHC panel for the diagnosis of hairy cell leukemia (HCL).

Application Notes

Optimal dilution of the recombinant ITGAE antibody should be determined by the researcher.

Immunogen

Recombinant human ITGAE partial protein was used as the immunogen for the recombinant ITGAE antibody.

Storage

Store the recombinant ITGAE antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).