
Keratin 14 Antibody KRT14/7977R / Cytokeratin 14 Rabbit MAb [clone KRT14/7977R] (V4492)

Catalog No.	Formulation	Size
V4492-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4492-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4492SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant RABBIT MONOCLONAL

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Rabbit
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG, kappa
Clone Name	KRT14/7977R
Purity	Protein A/G affinity
UniProt	P02533
Localization	Cytoplasm
Applications	Immunohistochemistry (FFPE): 1-2ug/ml for 30 minutes at RT
Limitations	This Keratin 14 Antibody KRT14/7977R is available for research use only.



IHC staining of FFPE human prostate tissue with Keratin 14 Antibody KRT14/7977R. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free recombinant Keratin 14 antibody (clone KRT14/7977R) as confirmation of integrity and purity.

Description

Keratin 14 antibody detects Cytokeratin 14, a member of the type I keratin family of intermediate filament proteins. It always pairs with the type II keratin K5 and form the primary keratin pair of the keratinocytes of stratified squamous epithelia, including the epidermis as well as mucosal non-keratinizing stratified squamous epithelia. Cytokeratin 14 is strongly expressed in the undifferentiated basal cell layer containing the stem cells and are down-regulated in the differentiating suprabasal cell layers. Otherwise, in the widely well stratified follicular outer root sheath, cytokeratin 14 is uniformly expressed throughout all layers. The expression spectrum of cytokeratin 14 in tumors corresponds well to the patterns in normal epithelia. Thus, most squamous cell carcinomas as well as malignant mesotheliomas strongly express this keratin whereas little, focal, or no expression is found in adenocarcinomas. Cytokeratin 14 may be a useful marker in the differential diagnosis of squamous cell carcinoma from other epithelial tumors. Recent studies also indicate that CK14 expression in breast cancer corresponded with poor clinical outcome and that CK14 may have diagnostic value in the sub-classification of NSCLC.

Explore our KRT14 Antibody KRT14/4128 for investigating cytokeratin 14 expression, basal epithelial differentiation, and intermediate filament organization in normal and diseased tissues.

Application Notes

Optimal dilution of the Keratin 14 Antibody KRT14/7977R should be determined by the researcher.

Immunogen

A recombinant human KRT14 fragment (within amino acids 272-472) was used as the immunogen for the Keratin 14 Antibody KRT14/7977R.

Storage

Aliquot the Keratin 14 Antibody KRT14/7977R and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.

Copyright © NSJ Bioreagents. All rights reserved.