


Recombinant Ki-67 Antibody [clone rMKI67/6499] (V8961)

| Catalog No. | Formulation | Size |
|----------------|---|--------|
| V8961-100UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 100 ug |
| V8961-20UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide | 20 ug |
| V8961SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free | 100 ug |


Recombinant **MOUSE MONOCLONAL**

[Bulk quote request](#)


| | |
|--------------------|---|
| Availability | 1-3 business days |
| Species Reactivity | Human |
| Format | Purified |
| Clonality | Recombinant Mouse Monoclonal |
| Isotype | Mouse IgG2a, kappa |
| Clone Name | rMKI67/6499 |
| Purity | Protein A/G affinity |
| UniProt | P46013 |
| Localization | Nuclear |
| Applications | Immunohistochemistry (FFPE) : 1-2ug/ml |
| Limitations | This recombinant Ki-67 antibody is available for research use only. |



IHC staining of FFPE human tonsil tissue with recombinant Ki67 antibody (clone rMKI67/6499). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



Negative control: IHC staining of FFPE human brain tissue using recombinant Ki-67 antibody (clone rMKI67/6499) at 2ug/ml in PBS for 30min RT. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free recombinant Ki-67 antibody (clone rMKI67/6499) as confirmation of integrity and purity.

Description

Ki-67 antigen is a nuclear, non-histone protein that is present in all stages of the cell cycle except G0. This characteristic makes Ki-67 an excellent marker for proliferating cells and is commonly used as one of the prognostic factors in cancer studies. A correlation has been demonstrated between Ki-67 index and the histo-pathological grade of neoplasms. Assessment of Ki-67 expression in renal and ureter tumors shows a correlation between tumor proliferation and disease progression, thus making it possible to differentiate high-risk patients. Ki-67 expression may also prove to be important for distinguishing between malignant and benign peripheral nerve sheath tumors. Ki-67 labeling index has been shown to be a prognostic marker in a number of neoplasms including grade II astrocytoma, oligodendroglioma, colon carcinoma, and breast carcinoma. In general, Ki-67 is a good marker of proliferating cell populations.

Application Notes

Optimal dilution of the recombinant Ki-67 antibody should be determined by the researcher.

Immunogen

A portion of amino acids 2293-2478 from the human protein was used as the immunogen for the recombinant Ki-67 antibody.

Storage

Aliquot the recombinant Ki-67 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.