
Recombinant MMP9 Antibody / Rabbit Monoclonal [clone MXMP9-1R] (V3755)

Catalog No.	Formulation	Size
V3755-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3755-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3755SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V3755IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml

Recombinant RABBIT MONOCLONAL

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Rabbit
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG, kappa
Clone Name	MXMP9-1R
Purity	Protein A affinity chromatography
UniProt	P14780
Localization	Cytoplasmic, nuclear, secreted
Applications	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT Prediluted IHC only format: incubate for 30 min at RT (1)
Limitations	This recombinant MMP9 antibody is available for research use only.



IHC testing of FFPE human spleen with recombinant MMP9 antibody (clone MXMP9-1R). Required HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min.

Description

Proteins of the matrix metalloproteinase (MMP) family are involved in the breakdown of extracellular matrix in normal physiological processes, such as embryonic development, reproduction, angiogenesis, bone development, wound healing, cell migration, learning and memory, as well as in pathological processes, such as arthritis, intracerebral hemorrhage, and metastasis. Most MMPs are secreted as inactive proproteins which are activated when cleaved by extracellular proteinases. The enzyme encoded by this gene degrades type IV and V collagens and other extracellular matrix proteins. Studies in rhesus monkeys suggest that the enzyme is involved in IL-8-induced mobilization of hematopoietic progenitor cells from bone marrow, and murine studies suggest a role in tumor-associated tissue remodeling. [Wiki]

Application Notes

The stated application concentrations are suggested starting points. Titration of the recombinant MMP9 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Amino acids 603-614 were used as the immunogen for the recombinant MMP9 antibody.

Storage

Store the recombinant MMP9 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).

Copyright © NSJ Bioreagents. All rights reserved.