


## Recombinant Mucin-1 Antibody / MUC1 [clone rMUC1/955] (V8513)

Catalog No.	Formulation	Size
V8513-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V8513-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V8513SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant **MOUSE MONOCLONAL**

**Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Recombinant Mouse Monoclonal
Isotype	Mouse IgG1, kappa
Clone Name	rMUC1/955
Purity	Protein G affinity chromatography
UniProt	P15941
Localization	Cytoplasmic, cell surface
Applications	Immunohistochemistry (FFPE) : 0.5-1ug/ml for 30 minutes at RT
Limitations	This recombinant Mucin-1 antibody is available for research use only.



IHC staining of FFPE human small intestine with recombinant Mucin-1 antibody. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free recombinant Mucin-1 antibody as confirmation of integrity and purity.

## Description

This MAb reacts with MUC1. The dominant epitope of this MAb has not yet been determined. MUC1 is a large cell surface mucin glycoprotein expressed by most glandular and ductal epithelial cells and some hematopoietic cell lineages. It is expressed on most secretory epithelium, including mammary gland and some hematopoietic cells. It is expressed abundantly in lactating mammary glands and over expressed abundantly in 90% breast carcinomas and metastases. Transgenic MUC1 has been shown to associate with all four c-erbB receptors and localize with c-erbB1 (EGFR) in lactating glands. The MUC1 gene contains seven exons and produces several different alternatively spliced variants. The major expressed form of MUC1 uses all seven exons and is a type 1 transmembrane protein with a large extracellular tandem repeat domain. The tandem repeat domain is highly O glycosylated and alterations in glycosylation have been shown in epithelial cancer cells. Antibody to EMA is useful as a pan-epithelial marker for detecting early metastatic loci of carcinoma in bone marrow or liver.

## Application Notes

Optimal dilution of the recombinant Mucin-1 antibody should be determined by the researcher.

## Immunogen

Recombinant full-length human MUC1 protein was used as the immunogen for the recombinant Mucin-1 antibody.

## Storage

Store the recombinant Mucin-1 antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).