


# Recombinant Napsin A Antibody [clone rNAPSA/1239] (V3634)

Catalog No.	Formulation	Size
V3634-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3634-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3634SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V3634IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml


## Recombinant MOUSE MONOCLONAL

### **Bulk quote request**


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Mouse Monoclonal
Isotype	Mouse IgG1, kappa
Clone Name	rNAPSA/1239
Purity	Protein G affinity chromatography
UniProt	O96009
Localization	Cytoplasmic
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT Western Blot : 2-4ug/ml
Limitations	This recombinant Napsin A antibody is available for research use only.



Western blot testing of human kidney tissue lysate with recombinant Napsin A antibody. Expected molecular weight: 38-45 kDa.



SDS-PAGE analysis of purified, BSA-free recombinant Napsin A antibody (clone rNAPSA/1239) as confirmation of integrity and purity.



IHC testing of FFPE human lung adenocarcinoma with recombinant Napsin A antibody (clone rNAPSA/1239). Required HIER: boil tissue sections in pH6, 10mM citrate buffer, for 10-20 min followed by cooling at RT for 20 min.

#### **Description**

Napsin-A is a protein that in humans is encoded by the NAPSA gene. The activation peptides of aspartic proteinases plays role as inhibitors of the active site. These peptide segments, or pro-parts, are deemed important for correct folding, targeting, and control of the activation of aspartic proteinase zymogens. The pronapsin A gene is expressed predominantly in lung and kidney. Its translation product is predicted to be a fully functional, glycosylated aspartic proteinase precursor containing an RGD motif and an additional 18 residues at its C-terminus. [Wiki]

#### **Application Notes**

Optimal dilution of the recombinant Napsin A antibody should be determined by the researcher.

1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

#### **Immunogen**

A recombinant fragment from the human protein (amino acids 189-299) was used as the immunogen for the recombinant Napsin A antibody.

#### **Storage**

Store the recombinant Napsin A antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).