
# Recombinant NKX2.2 Antibody / Rabbit Monoclonal [clone NX2/2198R] (V3668)

| Catalog No. | Formulation | Size |
|---|---|---|
| V3668-100UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide | 100 ug |
| V3668-20UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide | 20 ug |
| V3668SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free | 100 ug |
| V3668IHC-7ML | Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only* | 7 ml |

Recombinant RABBIT MONOCLONAL

**Bulk quote request**

| | |
|---|---|
| **Availability** | 1-3 business days |
| **Species Reactivity** | Human |
| **Format** | Purified |
| **Host** | Rabbit |
| **Clonality** | Recombinant Rabbit Monoclonal |
| **Isotype** | Rabbit IgG, kappa |
| **Clone Name** | NX2/2198R |
| **Purity** | Protein A affinity chromatography |
| **Gene ID** | 4821 |
| **Localization** | Nuclear |
| **Applications** | Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT |
| **Limitations** | This recombinant NKX2.2 antibody is available for research use only. |



IHC testing of FFPE Ewings sarcoma with recombinant NKX2.2 antibody (clone NX2/2198R). HIER: steam sections in 10mM citrate buffer, pH 6.0, for 10-20 min.

## Description

Expression of NKX2.2 has been found in neuroendocrine tumors of the gut, making it a potential marker for the study of gastrointestinal neuroendocrine tumors. More recently, NKX2.2 protein was identified as a target of EWS-FLI-1, the fusion protein specific to Ewing sarcoma, and was shown to be differentially upregulated in Ewing sarcoma on the basis of array-based gene expression analysis. Antibody to NKX2.2 detects a valuable marker for Ewing sarcoma, with a sensitivity of 93% and a specificity of 89%, and aids in the differential diagnosis of small round cell tumors.

## Application Notes

Titering of the recombinant NKX2.2 antibody may be required for optimal performance.

1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

## Immunogen

A human partial recombinant protein (within amino acids 1-119) was used as the immunogen for this recombinant NKX2.2 antibody.

## Storage

Store the recombinant NKX2.2 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).


Copyright © NSJ Bioreagents. All rights reserved.