


Recombinant NKX2.2 Antibody [clone rNX2/294] (V3756)

Catalog No.	Formulation	Size
V3756-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3756-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3756SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V3756IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml


Recombinant **MOUSE MONOCLONAL**

Bulk quote request

Species Reactivity	Human, Mouse, Rat
Format	Purified
Host	Mouse
Clonality	Recombinant Mouse Monoclonal
Isotype	Mouse IgG1, kappa
Clone Name	rNX2/294
Purity	Protein G affinity chromatography
Buffer	1X PBS, pH 7.4
UniProt	O95096
Gene ID	4821
Localization	Nuclear
Applications	Immunohistochemistry (FFPE) : 0.5-1ug/ml for 30 min at RT
Limitations	This recombinant NKX2.2 antibody is available for research use only.



IHC testing of Ewing's sarcoma stained with recombinant NKX2.2 antibody (clone rNX2/294). Required HIER: boil tissue sections in 10mM citrate buffer, pH 6, for 10-20 min followed by cooling at RT for 20 min.



SDS-PAGE analysis of purified, BSA-free recombinant NKX2.2 antibody (clone rNX2/294) as confirmation of integrity and purity.

Description

Expression of NKX2.2 has been found in neuroendocrine tumors of the gut, making it a potential marker for the study of gastrointestinal neuroendocrine tumors. More recently, NKX2.2 protein was identified as a target of EWS-FLI-1, the fusion protein specific to Ewing sarcoma, and was shown to be differentially upregulated in Ewing sarcoma on the basis of array-based gene expression analysis. Antibody to NKX2.2 detects a valuable marker for Ewing sarcoma, with a sensitivity of 93% and a specificity of 89%, and aids in the differential diagnosis of small round cell tumors.

Application Notes

The concentration stated for each application is a general starting point. Variations in protocols, secondaries and substrates may require the recombinant NKX2.2 antibody to be titrated up or down for optimal performance.

1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Full length human recombinant protein was used as the immunogen for this recombinant NKX2.2 antibody.

Storage

Store the recombinant NKX2.2 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).

References (2)