


Recombinant p16INK4a Antibody / CDKN2A [clone CDKN2A/4844R] (V9547)

Catalog No.	Formulation	Size
V9547-100UG	0.2~mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V9547-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V9547SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant RABBIT MONOCLONAL

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG, kappa
Clone Name	CDKN2A/4844R
Purity	Protein A/G affinity
UniProt	P42771
Localization	Nucleus, Cytoplasm
Applications	Western Blot : 1-2ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This recombinant p16INK4a antibody is available for research use only.



Western blot testing of human HepG2 cell lysate using recombinant p16INK4a antibody (clone CDKN2A/4844R). Predicted molecular weight ~16 kDa.



IHC staining of FFPE human cervical tissue with recombinant p16INK4a antibody (clone CDKN2A/4844R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human cervix with recombinant p16INK4a antibody (clone CDKN2A/4844R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

Description

p16INK4a is a tumor suppressor protein. It is a specific inhibitor of cdk4/cdk6, and a tumor suppressor involved in the pathogenesis of a variety of malignancies. Recent analyses of the p16INK4a gene revealed homozygous deletions, nonsense, missense, or frameshift mutations in several human cancers. Although the frequency of p16INK4a abnormalities is higher in tumor derived cell lines than in unselected primary tumors, significant subsets of clinical cases with aberrant p16INK4a gene have been reported among melanomas, gliomas, esophageal, pancreatic, lung, and urinary bladder carcinomas, and some types of leukemia. Expression of p16INK4a (p16 positive) is highly correlated with human papilloma virus (HPV) infection in head and neck squamous cell carcinomas (HNSCC). p16 status is an important prognostic indicator in HNSCC and the p16 positive/HPV16 negative group is likely a distinct subgroup lacking any HPV genotype.

Application Notes

Optimal dilution of the recombinant p16INK4a antibody should be determined by the researcher.

Immunogen

Recombinant full-length human p16INK4a protein was used as the immunogen for the recombinant p16INK4a antibody.

Storage

Aliquot the recombinant p16INK4a antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.