

Recombinant p63 Antibody [clone RM383] (R20340)

Catalog No.	Formulation	Size
R20340-0.1ML	Antibody in PBS with 50% glycerol, 1% BSA and 0.09% sodium azide	100 ul


Recombinant **RABBIT MONOCLONAL**

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG
Clone Name	RM383
Purity	Protein A purified from animal origin-free supernatant
UniProt	Q9H3D4
Gene ID	4288
Localization	Nuclear
Applications	Immunohistochemistry (FFPE) : 1:50-1:100 Western Blot : 1:500-1:1000
Limitations	This recombinant p63 antibody is available for research use only.

(kDa)
260-
160-
110-
80-
60-
50-
40-
30-
20-
15-
10-


Western blot test of human A431 cell lysate with recombinant p63 antibody at 1:1000.
Expected molecular weight: 63-77 kDa.



IHC testing of FFPE human tonsil tissue with recombinant p63 antibody. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 10-20 min and allow to cool before testing.



IHC testing of FFPE human breast tissue with recombinant p63 antibody. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 10-20 min and allow to cool before testing.



IHC testing of FFPE human prostate tissue with recombinant p63 antibody. HIER: boil tissue sections in pH6, 10mM citrate buffer, for 10-20 min and allow to cool before testing.

Description

This recombinant p63 antibody reacts to human p63.

Application Notes

The stated application concentrations are suggested starting points. Titration of the recombinant p63 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

A peptide corresponding to the C-terminal region of human p63 was used as the immunogen for this recombinant p63 antibody.

Storage

Store the recombinant p63 antibody at -20oC.