


Recombinant PCNA Antibody / Proliferating Cell Nuclear Antigen [clone rPCNA/8858] (V4825)

Catalog No.	Formulation	Size
V4825-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4825-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4825SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant MOUSE MONOCLONAL

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Mouse Monoclonal
Isotype	Mouse IgG1, kappa
Clone Name	rPCNA/8858
Purity	Protein A/G affinity
UniProt	P12004
Localization	Nucleus, Cytoplasm
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT Western Blot : 2-4ug/ml
Limitations	This recombinant PCNA antibody is available for research use only.




IHC staining of FFPE human tonsil tissue with recombinant PCNA antibody (clone rPCNA/8858) at 2ug/ml. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human tonsil tissue with recombinant PCNA antibody (clone rPCNA/8858) at 2ug/ml. Inset: PBS used in place of primary Ab (secondary Ab negative control). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



Western blot testing of human tumor cell lysates with recombinant PCNA antibody. Predicted molecular weight ~29 kDa, routinely observed at 29~36 kDa.



SDS-PAGE analysis of purified, BSA-free recombinant PCNA antibody (clone rPCNA/8858) as confirmation of integrity and purity.

Description

Recognizes a non-histone protein of 36kDa, which is identified as proliferating cell nuclear antigen (PCNA). It is also known as cyclin or polymerase delta auxiliary protein. Elevated expression of PCNA/cyclin has been shown in the nucleus during late G1 phase immediately before the onset of DNA synthesis, becoming maximal during S-phase and declining during G2 and M phases. This mAb is excellent for multiple applications.

Application Notes

Optimal dilution of the recombinant PCNA antibody should be determined by the researcher.

Immunogen

Recombinant full-length human PCNA protein was used as the immunogen for the recombinant PCNA antibody.

Storage

Aliquot the recombinant PCNA antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.

