


Recombinant Periostin Antibody / POSTN [clone POSTN/8523R] (V4324)

Catalog No.	Formulation	Size
V4324-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4324-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4324SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant **RABBIT MONOCLONAL**

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Rabbit
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG, kappa
Clone Name	POSTN/8523R
Purity	Protein A/G affinity
UniProt	Q15063
Localization	Secreted, Cytoplasm
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 minutes at RT
Limitations	This recombinant Periostin antibody is available for research use only.



Immunohistochemistry of Periostin antibody in human lung tissue. Formalin-fixed, paraffin-embedded human lung was stained with recombinant monoclonal Periostin antibody (clone POSTN/8523R) following heat-induced epitope retrieval by boiling in pH 9 10 mM Tris with 1 mM EDTA for 20 minutes and cooling prior to testing. HRP-DAB brown chromogenic signal highlights extracellular matrix and stromal regions along alveolar septa and peribronchial connective tissue, consistent with the secreted and matrix-associated localization of Periostin. The inset image shows PBS in place of the primary antibody as a secondary-only negative control, demonstrating minimal non-specific background staining.



IHC staining of FFPE human colon tissue with recombinant Periostin antibody (clone POSTN/8523R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free recombinant Periostin antibody (clone POSTN/8523R) as confirmation of integrity and purity.

Description

Periostin antibody recognizes Periostin, a secreted extracellular matrix protein encoded by the POSTN gene and also known as osteoblast specific factor 2. Recombinant Periostin Antibody (clone POSTN/8523R) is developed for research applications focused on detection of this matricellular protein in tissue sections and cell lysates. Periostin is synthesized with a signal peptide and secreted into the extracellular space, where it integrates into the interstitial matrix and basement membrane to regulate tissue structure and cell-matrix signaling.

Periostin antibody, also referred to as POSTN antibody and OSF-2 antibody, targets a member of the fasciclin family characterized by an N-terminal EMI domain and four tandem fasciclin-like domains. These domains enable interaction with integrins such as alpha v beta 3 and alpha v beta 5, as well as extracellular matrix components including collagen type I and fibronectin. Through these interactions, Periostin supports fibroblast activation, cytoskeletal organization, and collagen fibrillogenesis, particularly in mechanically stressed tissues.

POSTN expression is enriched in periosteum, periodontal ligament, heart valves, and fibrous connective tissues. During development and wound repair, Periostin is upregulated to promote matrix remodeling and tissue stabilization. In cardiovascular biology, it contributes to valvular morphogenesis and adaptive remodeling. Elevated expression has also been documented in fibrotic conditions, where it is associated with excessive extracellular matrix deposition and progressive tissue stiffening.

In oncology research, Periostin is frequently detected within tumor-associated stroma rather than malignant epithelial cells. Increased POSTN expression has been reported in breast, lung, colorectal, pancreatic, and ovarian cancers, often localized to cancer-associated fibroblasts and peritumoral connective tissue. This stromal pattern makes Periostin antibody valuable for investigation of tumor microenvironment dynamics, epithelial-mesenchymal transition, and metastatic progression.

The recombinant monoclonal clone POSTN/8523R provides specific recognition of Periostin for research use, supporting studies of extracellular matrix biology, fibrosis, development, and cancer-associated stromal remodeling at NSJ Bioreagents.

Application Notes

Optimal dilution of the recombinant Periostin antibody should be determined by the researcher.

Immunogen

A recombinant partial protein sequence (within amino acids 193-326) from the human protein was used as the immunogen for the Periostin antibody.

Storage

Aliquot the recombinant Periostin antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.