


# Recombinant PHH3 Antibody / Histone H3 (pSer10) [clone PHH3/471R] (V8647)

Catalog No.	Formulation	Size
V8647-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V8647-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V8647SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

## Recombinant RABBIT MONOCLONAL

### **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Rabbit Monoclonal
Isotype	Mouse IgG2a, kappa
Clone Name	PHH3/471R
Purity	Protein G affinity chromatography
UniProt	P68431
Localization	Nucleus
Applications	Immunohistochemistry (FFPE): 1-2ug/ml for 30 minutes at RT
Limitations	This recombinant PHH3 antibody is available for research use only.



IHC staining of FFPE human colon with recombinant PHH3 antibody. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

#### **Description**

Phosphohistone H3 (PHH3) is a marker specific for cells undergoing mitosis. Serine 10 of Histone H3 is phosphorylated

in association with mitotic chromatin condensation in late G2 and M phase of the cell cycle and thus, PHH3 can distinguish mitosis from apoptotic nuclei. The range of percentage PHH3 positive tumor nuclei was from 0.0 to 6.6% (median value 0.8%). Increased expression of PHH3 was significantly associated with tumor thickness (p = 0.031), presence of tumor ulceration (p = 0.041) and tumor necrosis (p = 0.027), but not with Clarks level of invasion. High levels of PHH3 was associated with increased mitotic count (p = 0.003) and high Ki-67 expression (p = 0.002). For central nervous system tumors, melanoma, soft tissue tumors, GIST, etc., PHH3 mAb is helpful for tumor pathological classification and prognosis.

#### **Application Notes**

Optimal dilution of the recombinant PHH3 antibody should be determined by the researcher.

#### **Immunogen**

A synthetic peptide corresponding to the amino acids surrounding phospho S10 were used as the immunogen for the recombinant PHH3 antibody.

#### **Storage**

Store the recombinant PHH3 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).