


Recombinant Rabbit IgG Fc Antibody [clone RMG02] (R20173)

Catalog No.	Formulation	Size
R20173-100UG	1 mg/ml in PBS with 50% glycerol, 1% BSA and 0.09% sodium azide	100 ug


Recombinant **GOAT MONOCLONAL**

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Rabbit
Format	Purified
Host	Goat
Clonality	Recombinant Goat Monoclonal
Isotype	Goat IgG
Clone Name	RMG02
Purity	Protein G purified from animal origin-free supernatant
Gene ID	N/A
Applications	ELISA : 0.01ug/ml-0.5ug/ml Western Blot (non-reduced) : 0.1-0.5ug/ml
Limitations	This recombinant Rabbit IgG Fc antibody is available for research use only.



Western blot of nonreduced(-) and reduced(+) rabbit IgG (20ng/lane), using 0.2ug/ml of the recombinant Rabbit IgG Fc antibody. This antibody reacts to nonreduced form (~150 kDa).



ELISA of IgGs from different species shows the recombinant Rabbit IgG Fc antibody reacts to the Fc region of rabbit IgG; no cross reactivity with human/mouse/rat IgG.



ELISA Titration: the plate was coated with different amounts of rabbit IgG. A serial dilution of the recombinant Rabbit IgG Fc antibody was used as the primary and an alkaline phosphatase conjugated anti-goat IgG as the secondary.

Description

This recombinant Rabbit IgG Fc antibody reacts to the Fc region of Rabbit IgG. No cross reactivity with human/mouse/rat IgG.

Application Notes

The stated application concentrations are suggested starting points. Titration of the recombinant Rabbit IgG Fc antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

Rabbit IgG was used as the immunogen for this recombinant Rabbit IgG Fc antibody.

Storage

Store the recombinant Rabbit IgG Fc antibody at -20oC (with glycerol) or aliquot and store at -20oC (without glycerol).