


Recombinant Rabbit IgG Fc Antibody [clone RMG02] (R20173)

| Catalog No. | Formulation | Size |
|--------------|---|--------|
| R20173-100UG | 1 mg/ml in PBS with 50% glycerol, 1% BSA and 0.09% sodium azide | 100 ug |


Recombinant **GOAT MONOCLONAL**

[Bulk quote request](#)


| | |
|--------------------|---|
| Availability | 1-3 business days |
| Species Reactivity | Rabbit |
| Format | Purified |
| Clonality | Recombinant Goat Monoclonal |
| Isotype | Goat IgG |
| Clone Name | RMG02 |
| Purity | Protein G purified from animal origin-free supernatant |
| Gene ID | N/A |
| Applications | ELISA : 0.01ug/ml-0.5ug/ml Western Blot (non-reduced) : 0.1-0.5ug/ml |
| Limitations | This recombinant Rabbit IgG Fc antibody is available for research use only. |



Western blot of nonreduced(-) and reduced(+) rabbit IgG (20ng/lane), using 0.2ug/ml of the recombinant Rabbit IgG Fc antibody. This antibody reacts to nonreduced form (~150 kDa).



ELISA of IgGs from different species shows the recombinant Rabbit IgG Fc antibody reacts to the Fc region of rabbit IgG; no cross reactivity with human/mouse/rat IgG.



ELISA Titration: the plate was coated with different amounts of rabbit IgG. A serial dilution of the recombinant Rabbit IgG Fc antibody was used as the primary and an alkaline phosphatase conjugated anti-goat IgG as the secondary.

Description

This recombinant Rabbit IgG Fc antibody reacts to the Fc region of Rabbit IgG. No cross reactivity with human/mouse/rat IgG.

Application Notes

The stated application concentrations are suggested starting points. Titration of the recombinant Rabbit IgG Fc antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

Rabbit IgG was used as the immunogen for this recombinant Rabbit IgG Fc antibody.

Storage

Store the recombinant Rabbit IgG Fc antibody at -20oC (with glycerol) or aliquot and store at -20oC (without glycerol).