


Recombinant RPA32 Antibody / Replication Protein A2 [clone RPA2/3140R] (V7886)

| Catalog No. | Formulation | Size |
|----------------|--|--------|
| V7886-100UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide | 100 ug |
| V7886-20UG | 0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide | 20 ug |
| V7886SAF-100UG | 1 mg/ml in 1X PBS; BSA free, sodium azide free | 100 ug |

Recombinant **RABBIT MONOCLONAL**

[Bulk quote request](#)

| | |
|--------------------|---|
| Availability | 1-3 business days |
| Species Reactivity | Human |
| Format | Purified |
| Host | Rabbit |
| Clonality | Recombinant Rabbit Monoclonal |
| Isotype | Rabbit IgG, kappa |
| Clone Name | RPA2/3140R |
| Purity | Protein A affinity chromatography |
| UniProt | P15927 |
| Applications | Western Blot : 1-2ug/ml |
| Limitations | This recombinant RPA32 antibody is available for research use only. |



Western blot testing of human MCF7 and T47D cell lysate with recombinant RPA32 antibody. Expected molecular weight ~32 kDa.

Description

Recognizes a protein of 34kDa, identified as a subunit of Replication Protein A (RPA) (also known as human single-

stranded DNA binding protein, or HSSB). RPA from human cells is a stable heterotrimer of 70kDa, 32-34kDa, and 11-14kDa subunits (RPA70, RPA32, and RPA14 respectively). It is involved in DNA replication, repair, and recombination. RPA is required for the SV40 large tumor antigen-catalyzed unwinding of SV40 DNA and stimulates DNA polymerase (pol) alpha and delta. RPA34 is phosphorylated at the G1/S boundary of the cell cycle or upon exposure of cells to DNA damage-inducing agents including ionizing and UV radiation.

Application Notes

Optimal dilution of the recombinant RPA32 antibody should be determined by the researcher.

Immunogen

A recombinant human full-length Replication protein A2 protein was used as the immunogen for this recombinant RPA32 antibody.

Storage

Store the recombinant RPA32 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).