


Recombinant Secretory Component Glycoprotein Antibody / ECM1 [clone rECM1/792] (V2484)

Catalog No.	Formulation	Size
V2484-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2484-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2484SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V2484IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml


Recombinant **MOUSE MONOCLONAL**

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Recombinant Mouse Monoclonal
Isotype	Mouse IgG1, kappa
Clone Name	rECM1/792
Purity	Protein G affinity chromatography
UniProt	Q16610
Localization	Cell surface and cytoplasmic
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT (1) Prediluted IHC Only Format : incubate for 30 min at RT (2)
Limitations	This recombinant Secretory Component Glycoprotein antibody is available for research use only.



IHC: Formalin-fixed, paraffin-embedded human colon carcinoma stained with recombinant Secretory Component Glycoprotein antibody (rECM1/792).



SDS-PAGE Analysis of Purified, BSA-Free recombinant Secretory Component Glycoprotein Antibody (clone rECM1/792). Confirmation of Integrity and Purity of the Antibody.

Description

This mAb reacts with a reduction-resistant epitope present in both free and SIgA bound Secretory Component. It does not react with the cell lines lacking secretory component. The antibody is useful for studying the distribution and level of both free and bound secretory component. Secretory component is differentially expressed in epithelium, and the antibody is a popular marker for identifying subpopulations of epithelial cells and epithelial differentiation. The Secretory component antibody is a useful research tool for studying mucosal immunity, inflammation, remodeling, differentiation and tumorigenesis, all processes associated with differential secretory component expression.

Application Notes

Optimal dilution of the recombinant Secretory Component Glycoprotein antibody should be determined by the researcher.

1. Staining of formalin-fixed tissues requires boiling tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min followed by cooling at RT for 20 min
2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Recombinant human protein was used as the immunogen for the recombinant Secretory Component Glycoprotein antibody.

Storage

Store the recombinant Secretory Component Glycoprotein antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).