


Recombinant SMMHC Antibody / Smooth Muscle Myosin Heavy Chain [clone MYH11/7087R] (V9683)

Catalog No.	Formulation	Size
V9683-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V9683-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V9683SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant **RABBIT MONOCLONAL**

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Rabbit
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG, kappa
Clone Name	MYH11/7087R
Purity	Protein A/G affinity
UniProt	P35749
Localization	Cytoplasm
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This recombinant SMMHC antibody is available for research use only.



IHC staining of FFPE human prostate tissue with recombinant SMMHC antibody MYH11/7087R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human smooth muscle tissue with recombinant SMMHC antibody (clone MYH11/7087R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free recombinant SMMHC antibody (clone MYH11/7087R) as confirmation of integrity and purity.

Description

Smooth Muscle Myosin, heavy chain (SMM-HC) is a cytoplasmic structural protein that is a major component of the contractile apparatus of the smooth muscle cells. Expression of smooth muscle myosin is developmentally regulated, appearing early and is specific for smooth muscle development. SM-MHC stains the intact myoepithelial cell (MEC) layers present in benign and in situ malignant breast and bronchioloalveolar lesions and is therefore very helpful in distinguishing between benign and malignant tumors. The antibody reacts with smooth muscle cells and myoepithelial cells, but not with myofibroblasts. It is very helpful in distinguishing between benign sclerosing breast lesions and infiltrating carcinomas in difficult cases since it strongly stains the myoepithelial layer in the benign lesions while it is negative in the infiltrating carcinomas.

Application Notes

Optimal dilution of the recombinant SMMHC antibody should be determined by the researcher.

Immunogen

A portion of amino acids 1-100 of Myosin protein was used as the immunogen for the recombinant SMMHC antibody.

Storage

Aliquot the recombinant SMMHC antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.