


# Recombinant STAT6 Antibody [clone STAT6/7163R] (V4052)

Catalog No.	Formulation	Size
V4052-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4052-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4052SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


## Recombinant RABBIT MONOCLONAL

## **Bulk quote request**


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG, kappa
Clone Name	STAT6/7163R
Purity	Protein A/G affinity
UniProt	P42226
Localization	Cytoplasm, nucleus
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This recombinant STAT6 antibody is available for research use only.



IHC staining of FFPE human renal cell carcinoma tissue with recombinant STAT6 antibody (clone STAT6/7163R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human liposarcoma tissue with recombinant STAT6 antibody (clone STAT6/7163R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human cerebellum tissue with recombinant STAT6 antibody (clone STAT6/7163R). Negative control inset: PBS used instead of primary antibody to control for secondary Ab binding. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

## **Description**

STAT6 is a transcription factor in the Jak/STAT signal transduction pathway responsible for mediating IL-4 immune signaling. STAT6 was recently suggested to be a reliable marker to distinguish solitary fibrous tumors from other soft tissue neoplasms. Gene fusions are common in solitary fibrous tumors. Recent next generation sequencing studies demonstrated the presence of a NAB2-STAT6 fusion, formed by an intrachromosomal inversion fusing two neighboring genes on chromosome 12q13, in 55-100% of solitary fibrous tumors, regardless of tumor morphology or anatomical site. By immunohistochemistry, nuclear STAT6 expression can discriminate solitary fibrous tumors from its morphological mimics in the meninges, including meningioma, glioblastoma, gliosarcoma, haemangioblastoma, schwannoma and haemangioma. A recent study by Cheah, et al. using the rabbit monoclonal STAT6 antibody (Clone YE361) observed expression in all solitary fibrous tumors (54/54) tested, regardless of histology, anatomical site or CD34 status. Morphological mimics of solitary fibrous tumors were negative, demonstrating 100% specificity.

### **Application Notes**

Optimal dilution of the recombinant STAT6 antibody should be determined by the researcher.

#### **Immunogen**

Recombinant full-length human protein was used as the immunogen for the recombinant STAT6 antibody.

#### **Storage**

Aliquot the recombinant STAT6 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.