


Recombinant SYP Antibody / Synaptophysin [clone rSYP/6856] (V9154)

Catalog No.	Formulation	Size
V9154-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V9154-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V9154SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant **MOUSE MONOCLONAL**

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Recombinant Mouse Monoclonal
Isotype	Mouse IgG1, kappa
Clone Name	rSYP/6856
Purity	Protein A/G affinity
UniProt	P08247
Localization	Cytoplasm
Applications	Immunohistochemistry (FFPE) : 2-4ug/ml
Limitations	This recombinant SYP antibody is available for research use only.



IHC staining of FFPE human brain tissue with recombinant SYP antibody (clone rSYP/6856). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free recombinant SYP antibody (clone rSYP/6856) as confirmation of integrity and purity.

Description

This recombinant mouse monoclonal antibody recognizes a protein of 38kDa that is identified as Synaptophysin. It is an N-glycosylated integral membrane protein found in neurons and endocrine cells. Synaptophysin contains four transmembrane domains and may function as a gap junction-like channel. This antibody identifies normal neuroendocrine cells and neuroendocrine neoplasms. Diffuse, finely granular, cytoplasmic staining is observed, which probably correlates with the distribution of the antigen within neurosecretory vesicles. Synaptophysin is an independent, broad-range marker of neural and neuroendocrine differentiation.

Application Notes

Optimal dilution of the recombinant SYP antibody should be determined by the researcher.

Immunogen

A portion of amino acids 224-313 was used as the immunogen for the recombinant SYP antibody.

Storage

Aliquot the recombinant SYP antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.