


## Recombinant TAG-72 Antibody [clone CA72/2869R] (V7779)

Catalog No.	Formulation	Size
V7779-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V7779-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V7779SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant **RABBIT MONOCLONAL**

[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Clonality</b>	Recombinant Rabbit Monoclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Name</b>	CA72/2869R
<b>Purity</b>	Protein A affinity chromatography
<b>Localization</b>	Cytoplasmic and cell surface
<b>Applications</b>	Immunohistochemistry (FFPE) : 1-2ug/ml
<b>Limitations</b>	This recombinant TAG-72 antibody is available for research use only.



IHC staining of human ovary with recombinant TAG-72 antibody (clone CA72/2869R).  
 HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 10-20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free recombinant TAG-72 antibody (clone CA72/2869R) as confirmation of integrity and purity.

## Description

TAG-72 is usually expressed by adenocarcinomas, but is negative in mesotheliomas. Studies have reported that this antibody has 80% sensitivity and 93% specificity for pulmonary adenocarcinoma. Therefore, TAG-72 is a useful marker to distinguish between mesothelioma and adenocarcinoma. However, false positive reactions can occur so results must be interpreted with the utmost caution. This antibody may be useful in the differentiation of non-small cell carcinomas from small cell carcinomas of the lung. The combined use of anti-TAG-72 and anti-GCDFP-15 is valuable in the diagnosis of apocrine carcinoma.

## Application Notes

Optimal dilution of the recombinant TAG-72 antibody should be determined by the researcher.

## Immunogen

Recombinant protein was used as the immunogen for the recombinant TAG-72 antibody.

## Storage

Store the recombinant TAG-72 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).