


TG Antibody Recombinant Mouse MAb r6E1 / Thyroglobulin Antibody [clone r6E1] (V3629)

Catalog No.	Formulation	Size
V3629-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3629-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3629SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V3629IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml


Recombinant **MOUSE MONOCLONAL**

[Bulk quote request](#)


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Recombinant Mouse Monoclonal
Isotype	Mouse IgG1, kappa
Clone Name	r6E1
Purity	Protein G affinity chromatography
UniProt	P01266
Localization	Cytoplasmic, secreted
Applications	Immunohistochemistry (FFPE) : 0.25-0.5ug/ml for 30 min at RT
Limitations	This recombinant TG antibody is available for research use only.



TG Antibody Recombinant Mouse MAb r6E1. Immunohistochemistry analysis of Thyroglobulin using TG Antibody Recombinant Mouse MAb r6E1 in FFPE human thyroid gland tissue section. HRP-DAB brown chromogenic staining highlights strong cytoplasmic signal in thyroid follicular epithelial cells forming thyroid follicles, with prominent staining of follicular luminal colloid material. The staining pattern is characteristic of Thyroglobulin expression in normal thyroid follicular epithelium and reflects its established role as a marker of thyroid follicular cell differentiation.



TG Antibody Recombinant Mouse MAb r6E1. Immunohistochemistry analysis of Thyroglobulin using TG Antibody Recombinant Mouse MAb r6E1 in FFPE human thyroid gland tissue section. HRP-DAB brown chromogenic staining highlights cytoplasmic signal in thyroid follicular epithelial cells forming well defined thyroid follicles, with strong staining of follicular luminal colloid material. The staining pattern is characteristic of Thyroglobulin expression in normal thyroid follicular epithelium and supports its role as a marker of thyroid follicular cell differentiation.



SDS-PAGE analysis of purified, BSA-free recombinant TG antibody (clone r6E1) as confirmation of integrity and purity.

Description

Thyroglobulin is a 660 kDa dimeric pre-protein with multiple glycosylation sites. It is produced by and processed within the thyroid gland to produce the hormone thyroxine and triiodothyronine. Prior to forming dimers, thyroglobulin monomers undergo conformational maturation in the endoplasmic reticulum. The vast majority of follicular carcinomas of the thyroid will give positive immunoreactivity for anti-thyroglobulin even though sometimes only focally. Poorly differentiated carcinomas of the thyroid are frequently anti-thyroglobulin negative. Adenocarcinomas of other-than-thyroid origin do not react with this antibody. This antibody is useful in identification of thyroid carcinoma of the papillary and follicular types. Presence of thyroglobulin in metastatic lesions establishes the thyroid origin of tumor. Anti-thyroglobulin, combined with anti-calcitonin, can identify medullary carcinomas of the thyroid. Furthermore, anti-thyroglobulin, combined with anti-TTF1, can be a reliable marker to differentiate between primary thyroid and lung neoplasms.

Application Notes

Optimal dilution of the TG Antibody Recombinant Mouse MAb r6E1 should be determined by the researcher.

1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Human thyroid follicular cells were used as the immunogen for the TG Antibody Recombinant Mouse MAb r6E1.

Storage

Store the recombinant TG antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).

Alternate Names

Thyroglobulin antibody, TG protein antibody, Thyroid hormone precursor protein antibody, Thyroid follicular cell marker antibody

