


Recombinant Thyroid Peroxidase Antibody [clone TPO/3813R] (V8137)

Catalog No.	Formulation	Size
V8137-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V8137-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V8137SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant **RABBIT MONOCLONAL**

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Rabbit
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG, kappa
Clone Name	TPO/3813R
Purity	Protein A affinity chromatography
UniProt	P07202
Localization	Cell surface, cytoplasmic
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This recombinant Thyroid Peroxidase antibody is available for research use only.



IHC staining of FFPE human thyroid carcinoma with recombinant Thyroid Peroxidase antibody (clone TPO/3813R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free recombinant Thyroid Peroxidase antibody (clone TPO/3813R) as confirmation of integrity and purity.

Description

Thyroperoxidase (TPO) is a 933-amino acid, type I transmembrane glycoprotein that plays a key role in thyroid gland function and autoimmunity. It is present as a dimer on the apical surface of thyroid follicular cells. TPO functions in the iodination of tyrosine residues in thyroglobulin and phenoxo-ester formation between pairs of iodinated tyrosines to generate the thyroid hormones, thyroxine and triiodothyronine. Mutations in this gene are associated with several disorders of thyroid hormonogenesis, including congenital hypothyroidism, congenital goiter, and thyroid hormone organification defect IIA. Malignant thyroid tumors exhibit an anomaly in TPO resulting in lower affinity for anti-TPO. This antibody may aid in the differentiation between benign and malignant thyroid tumors.

Application Notes

Optimal dilution of the recombinant Thyroid Peroxidase antibody should be determined by the researcher.

Immunogen

A recombinant human partial protein (amino acids 685-804) was used as the immunogen for this TPO antibody.

Storage

Store the recombinant Thyroid Peroxidase antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).