

Recombinant TP63 Antibody / Tumor protein 63 [clone rTP63/6941] (V9374)

Catalog No.	Formulation	Size
V9374-100UG	0.2~mg/ml in 1X PBS with $0.1~mg/ml$ BSA (US sourced), $0.05%$ sodium azide	100 ug
V9374-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V9374SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Recombinant MOUSE MONOCLONAL

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Mouse Monoclonal
Isotype	Mouse IgG2a, kappa
Clone Name	rTP63/6941
Purity	Protein A affinity
UniProt	Q9H3D4
Localization	Nucleus
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This recombinant TP63 antibody is available for research use only.



Description

p63 is a homolog of the tumor suppressor p53. It is identified in basal cells in the epithelial layers of a variety of tissues, including epidermis, cervix, urothelium, breast and prostate. p63 was detected in nuclei of the basal epithelium in normal prostate glands; however, it was not expressed in malignant tumors of the prostate. As a result, p63 has been reported as a useful marker for differentiating benign from malignant lesions in the prostate, particularly when used in combination

with markers of high molecular weight cytokeratins and the prostate-specific marker AMACR (P504S). p63 has also been shown to be a sensitive marker for lung squamous cell carcinomas (SqCC), with a sensitivity of ~90%. Specificity for lung SqCC, vs. lung adenocarcinoma (LADC), is approximately 80%. In breast tissue, p63 has been identified in myoepithelial cells of normal ducts.

Application Notes

Optimal dilution of the recombinant TP63 antibody should be determined by the researcher.

Immunogen

A portion of amino acids 1-200 was used as the immunogen for the recombinant TP63 antibody.

Storage

Aliquot the recombinant TP63 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.