


## Recombinant TPO Antibody / Thyroid Peroxidase [clone TPO/6417R] (V9282)

Catalog No.	Formulation	Size
V9282-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V9282-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V9282SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant **RABBIT MONOCLONAL**

[Bulk quote request](#)


<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Purified
<b>Host</b>	Rabbit
<b>Clonality</b>	Recombinant Rabbit Monoclonal
<b>Isotype</b>	Rabbit IgG
<b>Clone Name</b>	TPO/6417R
<b>Purity</b>	Protein A/G affinity
<b>UniProt</b>	P07202
<b>Localization</b>	Cytoplasm, Cell surface
<b>Applications</b>	Immunohistochemistry (FFPE) : 1-2ug/ml
<b>Limitations</b>	This recombinant TPO antibody is available for research use only.



IHC staining of FFPE human thyroid carcinoma tissue with recombinant TPO antibody (clone TPO/6417R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human thyroid carcinoma tissue with recombinant TPO antibody (clone TPO/6417R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free recombinant TPO antibody (clone TPO/6417R) as confirmation of integrity and purity.

## Description

Thyroperoxidase (TPO) is a 933 amino acid, type I transmembrane glycoprotein that plays a key role in thyroid gland function and autoimmunity. It is present as a dimer on the apical surface of thyroid follicular cells. TPO functions in the iodination of tyrosine residues in thyroglobulin and phenoxy-ester formation between pairs of iodinated tyrosines to generate the thyroid hormones, thyroxine and triiodothyronine. Mutations in this gene are associated with several disorders of thyroid hormonogenesis, including congenital hypothyroidism, congenital goiter, and thyroid hormone organification defect IIA. Malignant thyroid tumors exhibit an anomaly in TPO resulting in lower affinity for anti-TPO. This antibody may aid in the differentiation between benign and malignant thyroid tumors.

## Application Notes

Optimal dilution of the recombinant TPO antibody should be determined by the researcher.

## Immunogen

A portion of amino acids 685-804 from the human protein was used as the immunogen for the recombinant TPO antibody.

## Storage

Aliquot the recombinant TPO antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.