


Recombinant Tubulin Beta Antibody / TUBB3 [clone TUBB3/7090R] (V9479)

Catalog No.	Formulation	Size
V9479-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V9479-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V9479SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant **RABBIT MONOCLONAL**

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Rabbit
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG
Clone Name	TUBB3/7090R
Purity	Protein A/G affinity
UniProt	Q13509
Localization	Cytoplasm
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This recombinant Tubulin Beta antibody is available for research use only.



IHC staining of FFPE human ovarian carcinoma tissue with recombinant Tubulin Beta antibody (clone TUBB3/7090R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free recombinant Tubulin Beta antibody (clone TUBB3/7090R) as confirmation of integrity and purity.

Description

This gene encodes a class III member of the beta tubulin protein family. Beta tubulins are one of two core protein families (alpha and beta tubulins) that heterodimerize and assemble to form microtubules. This protein is primarily expressed in neurons and may be involved in neurogenesis and axon guidance and maintenance. Mutations in this gene are the cause of congenital fibrosis of the extraocular muscles type 3. In adults, tubulin beta 3 (TUBB3) is primarily expressed in neurons and is commonly used as a neuronal marker. It plays an important role in neuronal cell proliferation and differentiation.

Application Notes

Optimal dilution of the recombinant Tubulin Beta antibody should be determined by the researcher.

Immunogen

A portion of amino acids 437-450 from the human protein was used as the immunogen for the recombinant Tubulin Beta antibody.

Storage

Aliquot the recombinant Tubulin Beta antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.