


Recombinant TYR Antibody / Tyrosinase [clone rOCA1/812] (V3658)

Catalog No.	Formulation	Size
V3658-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3658-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3658SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V3658IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml


Recombinant **MOUSE MONOCLONAL**

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Recombinant Mouse Monoclonal
Isotype	Mouse IgG1, kappa
Clone Name	rOCA1/812
Purity	Protein G affinity chromatography
UniProt	P14679
Localization	Cytoplasmic
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml for 30 min at RT
Limitations	This recombinant Tyrosinase antibody is available for research use only.



IHC staining of FFPE human Melanoma with recombinant Tyrosinase antibody (clone rOCA1/812). Required HIER: boil tissue sections in 10mM Tris with 1mM EDTA, pH9, or 10mM citrate buffer, pH6, for 10-20 min followed by cooling at RT for 20 min.



SDS-PAGE analysis of purified, BSA-free recombinant Tyrosinase antibody (clone rOCA1/812) as confirmation of integrity and purity.

Description

This mAb shows no cross-reaction with MAGE-1 and tyrosinase-related protein 1, TRP-1/gp75. Tyrosinase is a copper-containing metalloglycoprotein that catalyzes several steps in the melanin pigment biosynthetic pathway; the hydroxylation of tyrosine to L-3,4-dihydroxy-phenylalanine (dopa), and the subsequent oxidation of dopa to dopaquinone. Mutations of the tyrosinase gene occur in various forms of albinism. Tyrosinase is one of the targets for cytotoxic T-cell recognition in melanoma patients. Staining of melanomas with this mAb shows tyrosinase in melanotic as well as amelanotic variants. This mAb is a useful marker for melanocytes and melanomas.

Application Notes

Optimal dilution of the recombinant Tyrosinase antibody should be determined by the researcher.

1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

Recombinant protein was used as the immunogen for this recombinant Tyrosinase antibody.

Storage

Store the recombinant Tyrosinase antibody at 2-8°C (with azide) or aliquot and store at -20°C or colder (without azide).