


Recombinant Ubiquitin Antibody [clone rUBB/9476] (V5804)

Catalog No.	Formulation	Size
V5804-100UG	0.2~mg/ml in 1X PBS with $0.1~mg/ml$ BSA (US sourced), $0.05%$ sodium azide	100 ug
V5804-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V5804SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant MOUSE MONOCLONAL

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Mouse Monoclonal
Isotype	Mouse IgG2b, kappa
Clone Name	rUBB/9476
Purity	Protein G affinity
UniProt	P62979, P62987
Localization	Cell membrane, cytoplasmic, nuclear
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml Western Blot : 2-4ug/ml
Limitations	This recombinant Ubiquitin antibody is available for research use only.



IHC staining of FFPE human brain tissue with recombinant Ubiquitin antibody (clone rUBB/9476). Inset: PBS used in place of primary Ab (secondary Ab negative control). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



Description

Ubiquitin is a highly conserved and plays an essential role in the ubiquitin-proteasome pathway. In ubiquitination process, it is first activated by forming a thiol-ester complex with the activation component E1, which is then transferred to ubiquitin-carrier protein E2, followed by to ubiquitin ligase E3 for final delivery to epsilon-NH2 of the target protein lysine residue. IkB, p53, cdc25A, Bcl-2 etc. are shown as targets of ubiquitin-proteasome process as part of regulation of cell cycle progression, differentiation, cell stress response, and apoptosis. Moreover, ubiquitin have been reported to bind covalently with pathological inclusions which are resistant to degradation e.g. neurofibrillary tangles/paired helical filaments in Alzheimer's disease, Lewy bodies seen in Parkinson's disease, and Pick bodies found in Pick's disease etc.

Application Notes

Optimal dilution of the recombinant Ubiquitin antibody should be determined by the researcher.

Immunogen

A recombinant partial protein from human Ubiquitin protein (within amino acids 1-119) was used as the immunogen for the recombinant Ubiquitin antibody.

Storage

Aliquot the recombinant Ubiquitin antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.