

Recombinant Villin Antibody [clone VIL1/4107R] (V8401)

Catalog No.	Formulation	Size
V8401-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V8401-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V8401SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Recombinant RABBIT MONOCLONAL

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG
Clone Name	VIL1/4107R
Purity	Protein A affinity chromatography
UniProt	P09327
Localization	Cytoplasmic & Cell surface
Applications	Immunohistochemistry (FFPE): 1-2ug/ml for 30 minutes at RT
Limitations	This recombinant Villin antibody is available for research use only.



IHC staining of FFPE human colon with recombinant Villin antibody (clone VIL1/4107R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human colon with recombinant Villin antibody (clone VIL1/4107R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

Description

Recognizes a protein of 95kDa, which is identified as villin. It is a major constituent in the microvilli, which compose the brush border of epithelial cells forming absorptive surfaces of the intestinal and renal proximal tubular epithelia. Anti-Villin labels the brush border area in the gastrointestinal mucosal epithelium and urogenital tract. Among neoplasms, villin is predominantly expressed in tumors of colorectal origin. Antibody to villin is useful in identifying malignant cells from primary and metastatic colorectal carcinomas. This antibody also labels Merkel cells of the skin.

Application Notes

Optimal dilution of the recombinant Villin antibody should be determined by the researcher.

Immunogen

A portion of amino acids 600-700 from the human protein was used as the immunogen for the recombinant Villin antibody.

Storage

Store the recombinant Villin antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).