


## Recombinant von Willebrand Factor Antibody / vWF [clone VWF/4384R] (V9636)

Catalog No.	Formulation	Size
V9636-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V9636-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V9636SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Recombinant **RABBIT MONOCLONAL**

[Bulk quote request](#)

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Rabbit
Clonality	Recombinant Rabbit Monoclonal
Isotype	Rabbit IgG
Clone Name	VWF/4384R
Purity	Protein A/G affinity
UniProt	P04275
Localization	Cytoplasm
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This recombinant von Willebrand Factor antibody is available for research use only.



IHC staining of FFPE human bone marrow with recombinant von Willebrand Factor antibody (clone VWF/4384R). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

## Description

Human von Willebrand factor (also called VWF and Factor VIII-related antigen) is a 309 kDa multimeric plasma

glycoprotein, important in the maintenance of hemostasis, it promotes adhesion of platelets to the sites of vascular injury by forming a molecular bridge between sub-endothelial collagen matrix and platelet-surface receptor complex GPIb-IX-V. Von Willebrand factor also acts as a chaperone for coagulation factor VIII, delivering it to the site of injury, stabilizing its heterodimeric structure and protecting it from premature clearance from plasma. Von Willebrand factor is synthesized by endothelial cells and is reported to be expressed in a number of tumors of vascular origin. IHC results aid in the classification of acute myeloid leukemia FAB type M7, angiosarcoma and epithelioid hemangioendothelioma.

## Application Notes

Optimal dilution of the recombinant von Willebrand Factor antibody should be determined by the researcher.

## Immunogen

A portion of amino acids 1150-1250 was used as the immunogen for the recombinant von Willebrand Factor antibody.

## Storage

Aliquot the recombinant von Willebrand Factor antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.