


Recombinant WT1 Antibody / Wilms Tumor 1 [clone rWT1/6908] (V9173)

Catalog No.	Formulation	Size
V9173-100UG	0.2~mg/ml in 1X PBS with $0.1~mg/ml$ BSA (US sourced), $0.05%$ sodium azide	100 ug
V9173-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V9173SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Recombinant MOUSE MONOCLONAL


Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Recombinant Mouse Monoclonal
Isotype	Mouse IgG1, kappa
Clone Name	rWT1/6908
Purity	Protein A/G affinity
UniProt	P19544
Localization	Nucleus and cytoplasm
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This recombinant WT1 antibody is available for research use only.



IHC staining of FFPE human ovarian carcinoma tissue with recombinant WT1 antibody (clone rWT1/6908). Negative control inset: PBS instead of primary antibody to control for secondary binding. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.

SDS-PAGE analysis of purified, BSA-free recombinant WT1 antibody (clone rWT1/6908) as confirmation of integrity and purity.



Description

The WT1 gene located at chromosome 11p13 codes for a transcription factor, a DNA-binding nucleoprotein, that plays a role primarily in the development of genitourinary organs. There are at least eight isoforms ranging between 52 and 62kDa produced by a combination of alternative splicing and RNA editing. WT1 is synthesized and reside in the cytoplasm in an inactive form. When activated through phosphorylation it is translocated to the nucleus. In tumor tissues, WT1 is detected in tumor cells of Wilms Tumor (also known as nephroblastoma) and mesothelioma. Additionally, WT1 expression has been found in ovarian serous carcinomas and some breast carcinomas. WT1 is particularly used for distinguishing malignant mesothelioma and ovarian serous carcinoma from non-serous carcinomas.

Application Notes

Optimal dilution of the recombinant WT1 antibody should be determined by the researcher.

Immunogen

A portion of amino acids 1-100 was used as the immunogen for the recombinant WT1 antibody.

Storage

Aliquot the recombinant WT1 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.