


# Renalase Antibody [clone RNLS/1940] (V3904)


Catalog No.	Formulation	Size
V3904-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V3904-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V3904SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

## **Bulk quote request**

Availability	1-2 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b, kappa
Clone Name	RNLS/1940
Purity	Protein G affinity
UniProt	Q5VYX0
Localization	Secreted
Applications	ELISA (order BSA/sodium Azide-free Format For Coating) :
Limitations	This Renalase antibody is available for research use only.



IHC testing of FFPE human kidney with Renalase antibody (clone RNLS/1940). HIER: boil tissue sections in pH6, 10mM citrate buffer, for 10-20 min followed by cooling at RT for 20 min.



IHC testing of FFPE human kidney with Renalase antibody (clone RNLS/1940). HIER: boil tissue sections in pH6, 10mM citrate buffer, for 10-20 min followed by cooling at RT for 20 min.

### **Description**

Renalase is a 342 amino acid FAD-dependent amine oxidase that is highly expressed in kidney and is expressed at a lower level in heart, skeletal muscle and small intestine. Renalase is secreted in the blood by the kidney and it is thought to regulate cardiac function and systemic blood pressure. It is also suggested that Renalase functions as a hormone that metabolizes circulating catecholamines, which have an active role in the sympathetic and parasympathetic nervous systems. Individuals with chronic kidney disease and end-stage renal disease have markedly reduced levels of plasma Renalase than healthy individuals. Infusion of Renalase in animal models causes decrease in heart rate, cardiac contractility and blood pressure. Two isoforms of Renalase exist due to alternative splicing events.

### **Application Notes**

Optimal dilution of the Renalase antibody should be determined by the researcher.

### **Immunogen**

A portion of amino acids 34-235 from the human protein was used as the immunogen for the Renalase antibody.

### **Storage**

Store the Renalase antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).