

## RICTOR Antibody (F53372)

Catalog No.	Formulation	Size
F53372-0.4ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.4 ml
F53372-0.08ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.08 ml

[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human
<b>Format</b>	Antigen affinity purified
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit Ig
<b>Purity</b>	Antigen affinity
<b>UniProt</b>	Q6R327
<b>Applications</b>	Western Blot : 1:2000
<b>Limitations</b>	This RICTOR antibody is available for research use only.



Western blot testing of RICTOR antibody at 1:2000 dilution + HepG2 lysate; Predicted molecular weight ~200 kDa.

## Description

Subunit of mTORC2, which regulates cell growth and survival in response to hormonal signals. mTORC2 is activated by growth factors, but, in contrast to mTORC1, seems to be nutrient- insensitive. mTORC2 seems to function upstream of Rho GTPases to regulate the actin cytoskeleton, probably by activating one or more Rho-type guanine nucleotide exchange factors. mTORC2 promotes the serum-induced formation of stress-fibers or F-actin. mTORC2 plays a critical role in AKT1 'Ser-473' phosphorylation, which may facilitate the phosphorylation of the activation loop of AKT1 on 'Thr-308' by PDK1 which is a prerequisite for full activation. mTORC2 regulates the phosphorylation of SGK1 at 'Ser-422'. mTORC2 also modulates the phosphorylation of PRKCA on 'Ser-657'. Plays an essential role in embryonic growth and development.

## Application Notes

Titration of the RICTOR antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

This RICTOR antibody was produced from a rabbit immunized with a KLH conjugated synthetic peptide between 1068-1102 amino acids from the Central region of human RICTOR.

## Storage

Aliquot the RICTOR antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.