


RING2 Antibody / RNF2 / RING1B (RQ7199)


Catalog No.	Formulation	Size
RQ7199	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Antigen affinity purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	Q99496
Localization	Cytoplasmic, nuclear
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 2-5ug/ml Direct ELISA : 0.1-0.5ug/ml
Limitations	This RING2 antibody is available for research use only.




IHC staining of FFPE human laryngeal squamous carcinoma tissue with RING2 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human thyroid cancer tissue with RING2 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human placental tissue with RING2 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Western blot testing of human 1) HL60, 2) K562, 3) U-2 OS and 3) Caco-2 cell lysate with RING2 antibody. Predicted molecular weight ~38 kDa.

Description

E3 ubiquitin-protein ligase RING2 is an enzyme that in humans is encoded by the RNF2 gene. Polycomb group (PcG) of proteins form the multiprotein complexes that are important for the transcription repression of various genes involved in development and cell proliferation. The protein encoded by this gene is one of the PcG proteins. It has been shown to interact with, and suppress the activity of, transcription factor CP2 (TFCP2/CP2). Studies of the mouse counterpart suggested the involvement of this gene in the specification of anterior-posterior axis, as well as in cell proliferation in early development. This protein was also found to interact with huntingtin interacting protein 2 (HIP2), an ubiquitin-conjugating enzyme, and possess ubiquitin ligase activity.

Application Notes

Optimal dilution of the RING2 antibody should be determined by the researcher.

Immunogen

Recombinant human protein (amino acids M1-K336) was used as the immunogen for the RING2 antibody.

Storage

After reconstitution, the RING2 antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.

