


## RIP2 Antibody (R30960)

Catalog No.	Formulation	Size
R30960	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

[Bulk quote request](#)

<b>Availability</b>	1-3 business days
<b>Species Reactivity</b>	Human, Mouse, Rat
<b>Format</b>	Antigen affinity purified
<b>Clonality</b>	Polyclonal (rabbit origin)
<b>Isotype</b>	Rabbit IgG
<b>Purity</b>	Antigen affinity
<b>Buffer</b>	Lyophilized from 1X PBS with 2.5% BSA and 0.025% sodium azide/thimerosal
<b>UniProt</b>	O43353
<b>Applications</b>	Western Blot : 0.5-1ug/ml
<b>Limitations</b>	This RIP2 antibody is available for research use only.



Western blot testing of RIP2 antibody and Lane 1: A549; 2: HeLa; 3: PANC; 4: COLO320 cell lysate. Predicted molecular weight: ~61 kDa.

## Description

Receptor-interacting serine/threonine-protein kinase 2, also known as CARD3, CARDIAK, RICK, and RIP2, is an enzyme that in humans is encoded by the RIPK2 gene. It has 540-amino acid protein in length. Northern blot analysis revealed that RIPK2 is expressed in various human tissues as 2.5- and 1.8-kb mRNAs that differ due to alternative polyadenylation. It is a novel kinase that may regulate apoptosis induced by the FAS receptor pathway. This gene encodes a member of the receptor-interacting protein (RIP) family of serine/threonine protein kinases. The encoded protein contains a C-terminalcaspase recruitment domain(CARD), and is a component of signaling complexes in both the innate and adaptive immune pathways. It is a potent activator of NF-kappa B and inducer of apoptosis in response to various stimuli, CARDIAK (CARD-containing ICE-associated kinase) specifically interacted with the CARD of

ICE/caspase-1, and this interaction correlated with the processing of pro-caspase-1 and the formation of the active caspase-1 p20 subunit.

## Application Notes

The stated application concentrations are suggested starting amounts. Titration of the RIP2 antibody may be required due to differences in protocols and secondary/substrate sensitivity.

## Immunogen

An amino acid sequence from the C-terminus of human RIP2 (DIQGEEFAKVIVQKLKDNKQ) was used as the immunogen for this RIP2 antibody.

## Storage

After reconstitution, the RIP2 antibody can be stored for up to one month at 4°C. For long-term, aliquot and store at -20°C. Avoid repeated freezing and thawing.