

RPAP3 Antibody / RNA polymerase II-associated protein 3 (FY12317)

Catalog No.	Formulation	Size
FY12317	Adding 0.2 ml of distilled water will yield a concentration of 500 ug/ml	100 ug

Bulk quote request

Availability	1-2 days
Species Reactivity	Human, Mouse, Rat
Format	Lyophilized
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Immunogen affinity purified
Buffer	Each vial contains 4 mg Trehalose, 0.9 mg NaCl, 0.2 mg Na2HPO4.
UniProt	Q9H6T3
Applications	Western Blot: 0.25-0.5ug/ml ELISA: 0.1-0.5ug/ml
Limitations	This RPAP3 antibody is available for research use only.

Description

RPAP3 antibody detects RNA polymerase II-associated protein 3, encoded by the RPAP3 gene on chromosome 12q24.31. RPAP3 antibody is commonly used in transcription, chaperone, and cancer research. RPAP3 is a component of the R2TP cochaperone complex, which cooperates with HSP90 and HSP70 to stabilize and assemble large macromolecular complexes. It is involved in the maturation of RNA polymerase II, snoRNPs, PIKKs, and other critical protein assemblies.

Structurally, RPAP3 is a ~75 kDa protein with multiple tetratricopeptide repeat (TPR) domains, which mediate protein-protein interactions. These domains allow RPAP3 to bridge HSP90 and PIH1D1, central elements of the R2TP complex. Its N-terminal domain anchors it to the cochaperone machinery, while its C-terminal region facilitates substrate recognition and client protein assembly.

Functionally, RPAP3 stabilizes RNA polymerase II subunits, enabling proper transcriptional initiation and elongation. It also regulates assembly of snoRNPs, which support ribosomal RNA modification, and PIKKs, which are central to DNA damage response, nutrient sensing, and stress signaling. Researchers use RPAP3 antibody to explore protein complex maturation, transcriptional regulation, and chaperone pathways.

Clinically, RPAP3 dysregulation has been linked to cancer progression. Because it stabilizes oncogenic PIKKs such as mTOR, ATR, and ATM, RPAP3 supports tumor growth and therapy resistance. Its expression patterns may serve as biomarkers for prognosis and treatment response. NSJ Bioreagents provides RPAP3 antibody to support transcription, DNA damage response, and cancer research.

Experimentally, RPAP3 antibody is applied in western blotting to detect the ~75 kDa protein, in immunoprecipitation to study R2TP complex partners, and in immunofluorescence microscopy to visualize nuclear and cytoplasmic distribution. Chromatin immunoprecipitation using RPAP3 antibody helps examine transcriptional regulation via RNA polymerase II stabilization.

Application Notes

Optimal dilution of the RPAP3 antibody should be determined by the researcher.

Immunogen

E.coli-derived human RPAP3 recombinant protein (Position: R84-E604) was used as the immunogen for the RPAP3 antibody.

Storage

After reconstitution, the RPAP3 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.