



RXR gamma Antibody / RXRG [clone PCRP-RXRG-5C9] (V4753)

Catalog No.	Formulation	Size
V4753-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4753-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4753SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug


Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b
Clone Name	PCRP-RXRG-5C9
Purity	Protein A/G affinity
UniProt	P78347
Localization	Nucleus
Applications	Flow Cytometry : 1-2ug/million cells
Limitations	This RXR gamma antibody is available for research use only.





Flow cytometry testing of PFA-fixed human HeLa cells with RXR gamma antibody (clone PCRP-RXRG-5C9) followed by goat anti-mouse IgG-CF488 (blue); Red = unstained cells.



SDS-PAGE analysis of purified, BSA-free RXR gamma antibody (clone PCRP-RXRG-5C9) as confirmation of integrity and purity.

Description

Two families of retinoid receptors, RARs and RXRs, have been identified. Retinoic acid receptors (RARs) include RAR α , RAR β and RAR γ , each of which have a high affinity for all trans-retinoic acids and belong to the same class of nuclear transcription factors as thyroid hormone receptors, vitamin D 3 receptor and ecdysone receptor. The ligand-binding domains of the RARs are highly conserved and RAR isoforms are expressed in distinct patterns throughout development and in the mature organism. Members of the retinoid X receptor (RXR) family, RXR α , RXR β and RXR γ , are activated by 9-cis-RA, a stereo- and photo-isomer of all trans-RA that is expressed in vivo in both liver and kidney and may represent a widely used hormone. As is true for the RAR subfamily, the RXR receptors are closely related to each other both in their DNA-binding and ligand-binding domains and are encoded by separate genes at distinct chromosomal loci.

Application Notes

Optimal dilution of the RXR gamma antibody should be determined by the researcher.

Immunogen

Recombinant human RXR gamma protein was used as the immunogen for the RXRG antibody.

Storage

Aliquot the RXR gamma antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.