# SARS-CoV-2 NSP8 Antibody (RQ6303)

| Catalog No. | Formulation | Size |
|---|---|---|
| RQ6303 | 0.5mg/ml if reconstituted with 0.2ml sterile DI water | 100 ug |

**Bulk quote request**

| | |
|---|---|
| **Availability** | 1-3 business days |
| **Species Reactivity** | Human |
| **Format** | Antigen affinity purified |
| **Host** | Rabbit |
| **Clonality** | Polyclonal (rabbit origin) |
| **Isotype** | Rabbit IgG |
| **Purity** | Affinity purified |
| **Buffer** | Lyophilized from 1X PBS with 2% Trehalose |
| **UniProt** | P0DTC1/P0DTD1 |
| **Applications** | ELISA |
| **Limitations** | This SARS-CoV-2 NSP8 antibody is available for research use only. |



## Description

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is an enveloped, positive-sense, single-stranded RNA virus that causes coronavirus disease 2019 (COVID-19). Virus particles include the RNA genetic material and structural proteins needed for invasion of host cells. Once inside the cell the infecting RNA is used to encode structural proteins that make up virus particles, nonstructural proteins that direct virus assembly, transcription, replication and host control and accessory proteins whose function has not been determined.~ ORF1ab, the largest gene, contains overlapping open reading frames that encode polyproteins PP1ab and PP1a. The polyproteins are cleaved to yield 16 nonstructural proteins, NSP1-16. Production of the longer (PP1ab) or shorter protein (PP1a) depends on a -1 ribosomal frameshifting

event. The proteins, based on similarity to other coronaviruses, include the papain-like proteinase protein (NSP3), 3C-like proteinase (NSP5), RNA-dependent RNA polymerase (NSP12, RdRp), helicase (NSP13, HEL), endoRNAse (NSP15), 2'-O-Ribose-Methyltransferase (NSP16) and other nonstructural proteins. SARS-CoV-2 nonstructural proteins are responsible for viral transcription, replication, proteolytic processing, suppression of host immune responses and suppression of host gene expression. The RNA-dependent RNA polymerase is a target of antiviral therapies.

## Application Notes

Optimal dilution of the SARS-CoV-2 NSP8 antibody should be determined by the researcher.

## Immunogen

Amino acids AIASEFSSLPSYAAFATAQEAYEQAVANGDSEVVLKKLKKSLNVAKSEFDRDAAMQRKLEKMADQAMTQMY KQARSEDKRAKVTSAMQTMLFTMLRKLDNDALNNIINNARD GCVPLNIIPLTTAAKLMVVIPDYNTYKNTCDGTTFTYASALWEIQQVVDADSKIVQLSEISMDNSPNLAWPLIVTALRANSAV KLQ were used as the immunogen for the SARS-CoV-2 NSP8 antibody.

## Storage

After reconstitution, the SARS-CoV-2 NSP8 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.


Copyright © NSJ Bioreagents. All rights reserved.