SATB2 Antibody [clone rSATB2/8635] (V4930)

Catalog No.	Formulation	Size
V4930-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V4930-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V4930SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Recombinant MOUSE MONOCLONAL

Bulk quote request

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Recombinant Mouse Monoclonal
Isotype	Mouse IgG1, kappa
Clone Name	rSATB2/8635
Purity	Protein A/G affinity
UniProt	Q9UPW6
Localization	Nucleus
Applications	ELISA (Order BSA-free format for coating)
Limitations	This SATB2 antibody is available for research use only.



SDS-PAGE analysis of purified, BSA-free SATB2 antibody (clone rSATB2/8635) as confirmation of integrity and purity.

Description

SATB2 is a DNA binding protein that specifically binds nuclear matrix attachment regions. It is involved in transcription regulation and chromatin remodeling. SATB2 expression in colorectal carcinomas (CRC) is correlated with good

prognosis and in laryngeal squamous cell carcinoma it functions as a tumor suppressor wherein loss of expression is positively correlated with high tumor grade and recurrence. Moreover, SATB2, in combination with CK20, could identify almost all CRC s. Upper gastrointestinal (GI) carcinomas and pancreatic ductal carcinomas are usually negative for SATB2, and ovarian carcinomas, lung adenocarcinomas, and adenocarcinomas from other origin are rarely positive for SATB2. Additionally, SATB2 antibody can identify neuroendocrine neoplasms of colon and rectum because SATB2 is usually negative in neuroendocrine neoplasms of the GI tract, pancreas, and lung. More recently, it has been reported that SATB2 is a sensitive marker for tumors with osteoblastic differentiation.

Researchers investigating colorectal tumor markers, epithelial lineage determination, and chromatin-associated transcriptional regulation may also be interested in our SATB2 Antibody / Colorectal and Lineage Marker page featuring validated immunohistochemistry and western blot applications for colorectal pathology research.

Application Notes

Optimal dilution of the SATB2 antibody should be determined by the researcher.

Immunogen

A recombinant partial protein sequence (within amino acids 150-350) from the human protein was used as the immunogen for the SATB2 antibody.

Storage

Aliquot the SATB2 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.

Copyright © NSJ Bioreagents. All rights reserved.