


SBDS Antibody / Shwachman-Bodian-Diamond syndrome protein (RQ7423)


| Catalog No. | Formulation | Size |
|-------------|---|--------|
| RQ7423 | 0.5mg/ml if reconstituted with 0.2ml sterile DI water | 100 ug |

Bulk quote request


| Availability | 1-3 business days |
|--------------------|--|
| Species Reactivity | Human, Mouse, Rat |
| Format | Antigen affinity purified |
| Clonality | Polyclonal (rabbit origin) |
| Isotype | Rabbit IgG |
| Purity | Antigen affinity purified |
| Buffer | Lyophilized from 1X PBS with 2% Trehalose |
| UniProt | Q9Y3A5 |
| Localization | Nuclear, cytoplasmic |
| Applications | Western Blot : 0.5-1ug/ml Immunofluorescence : 5ug/ml Flow Cytometry : 1-3ug/million cells Direct ELISA : 0.1-0.5ug/ml |
| Limitations | This SBDS antibody is available for research use only. |



Immunofluorescent staining of FFPE human U-2 OS cells with SBDS antibody (green) and Beta Tubulin mAb (red). HIER: steam section in pH6 citrate buffer for 20 min.



Western blot testing of 1) human 293T, 2) human HepG2, 3) human HL60, 4) human SH-SY5Y, 5) rat brain, 6) rat lung, 7) mouse brain and 8) mouse lung tissue lysate with SBDS antibody. Predicted molecular weight ~29 kDa.



Flow cytometry testing of fixed and permeabilized human HEL cells with SBDS antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= SBDS antibody.

Description

Ribosome maturation protein SBDS is a protein that in humans is encoded by the SBDS gene. This gene encodes a highly conserved protein that plays an essential role in ribosome biogenesis. The encoded protein interacts with elongation factor-like GTPase 1 to disassociate eukaryotic initiation factor 6 from the late cytoplasmic pre-60S ribosomal subunit allowing assembly of the 80S subunit. Mutations within this gene are associated with the autosomal recessive disorder Shwachman-Bodian-Diamond syndrome. This gene has a closely linked pseudogene that is distally located.

Application Notes

Optimal dilution of the SBDS antibody should be determined by the researcher.

Immunogen

E. coli-derived recombinant human protein (amino acids R19-D241) was used as the immunogen for the SBDS antibody.

Storage

After reconstitution, the SBDS antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.