


SELL Antibody / L-Selectin / LAM-1 (F55079)


Catalog No.	Formulation	Size
F55079-0.2ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.2 ml
F55079-0.05ML	In 1X PBS, pH 7.4, with 0.09% sodium azide	0.05 ml

[Bulk quote request](#)


Availability	1-2 business days
Species Reactivity	Human
Format	Purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit Ig
UniProt	P14151
Applications	Western Blot : 1:1000-1:2000 Immunohistochemistry (FFPE) : 1:10-1:50 Flow Cytometry : 1:25 per million cells in 0.1ml
Limitations	This SELL antibody is available for research use only.



IHC staining of FFPE human lymph tissue with SELL antibody. HIER: steam section in pH6 citrate buffer for 20 min and allow to cool prior to staining.



Western blot testing of human 1) Raji, 2) Jurkat, 3) K562, 4) MCF7 and 5) HL60 cell lysate with SELL antibody. Expected/observed molecular weight: 42~95 kDa depending on level of glycosylation.



Description

L-Selectin, also known as CD62L, SELL and LAM-1, is a cell adhesion molecule found on the surface of leukocytes, or white blood cells. It plays a role in the immune response by facilitating the binding of leukocytes to endothelial cells in the blood vessels, allowing them to migrate to sites of infection or inflammation. One of the key functions of L-Selectin is its involvement in the process of leukocyte rolling. When an infection or injury occurs, inflammatory signals are released that trigger the expression of adhesion molecules on the surface of endothelial cells. L-Selectin on leukocytes then binds to its ligands on the endothelial cells, causing the leukocytes to roll along the vessel wall. This rolling motion slows down the leukocytes and allows them to survey the endothelial surface for signs of infection. Once the leukocytes have identified the site of infection, they can firmly adhere to the endothelial cells and extravasate, or migrate, into the surrounding tissue. L-Selectin plays a critical role in this step by mediating the initial interaction between leukocytes and endothelial cells. Without L-Selectin, leukocytes would not be able to efficiently migrate to the site of infection, impairing the immune response.

In addition to its role in leukocyte trafficking, L-Selectin also plays a role in regulating the activation and function of leukocytes. Studies have shown that L-Selectin signaling can modulate the production of cytokines and chemokines, which are key mediators of inflammation. By influencing the inflammatory response, L-Selectin can help regulate the magnitude and duration of immune reactions, preventing excessive tissue damage. Furthermore, L-Selectin has been implicated in a variety of inflammatory and autoimmune diseases. Dysregulation of L-Selectin expression or function can lead to abnormal leukocyte trafficking and contribute to chronic inflammation. Understanding the role of L-Selectin in these conditions may provide insights into potential therapeutic targets for treating inflammatory diseases.

Application Notes

Titration of the SELL antibody may be required due to differences in protocols and secondary/substrate sensitivity.

Immunogen

A portion of amino acids 346-372 from the human protein was used as the immunogen for this SELL antibody.

Storage

Aliquot the SELL antibody and store frozen at -20°C or colder. Avoid repeated freeze-thaw cycles.