
SERBP1 Antibody / PAI-RBP1 [clone SERBP1/3493] (V7599)


Catalog No.	Formulation	Size
V7599-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V7599-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V7599SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V7599IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Host	Mouse
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b, kappa
Clone Name	SERBP1/3493
Purity	Protein G affinity chromatography
UniProt	Q8NC51
Localization	Cytoplasmic
Applications	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT Western blot: 1-2ug/ml
Limitations	This SERBP1 antibody is available for research use only.




IHC testing of FFPE human urothelial carcinoma with SERBP1 antibody (clone SERBP1/3493). Required HIER: boil tissue sections in pH6, 10mM citrate buffer, for 10-20 min followed by cooling at RT for 20 min.




IHC testing of FFPE human urothelial carcinoma with SERBP1 antibody (clone SERBP1/3493). Required HIER: boil tissue sections in pH6, 10mM citrate buffer, for 10-20 min followed by cooling at RT for 20 min.



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using SERBP1 antibody (clone SERBP1/3493). These results demonstrate the foremost specificity of the SERBP1/3493 mAb.
Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.



SDS-PAGE analysis of purified, BSA-free SERBP1 antibody (clone SERBP1/3493) as confirmation of integrity and purity.



Western blot testing of human 1) K562 and 2) PC-3 cell lysate with SERBP1 antibody. Predicted molecular weight ~45 kDa but observed at 45-60 kDa.

Description

PAI-RBP1 (plasminogen activator inhibitor 1 RNA-binding protein), also known as SERBP1 (SERPINE1 mRNA-binding protein 1), CGI-55, CHD3IP (chromodomain helicase DNA binding protein 3 interacting protein), HABP4L or PAI-RBP1, is a membrane-associated protein that localizes to the nucleus, the perinuclear region of the cytoplasm and the plasma membrane. PAI-RBP1 is believed to play a role in the regulation of mRNA stability, as it specifically binds to the CRS (cyclic nucleotide-responsive sequence) motif of the PAI-1 mRNA and acts to stabilize the mRNA and regulate its expression. In addition, PAI-RBP1 interacts with Mi2-alpha and may be involved in chromatin remodeling. PAI-RBP1 also interacts with PGRMC1 and participates in the transduction of Progesterone's antiapoptotic action in ovarian cell types. The gene encoding PAI-RBP1 is overexpressed in ovarian cancer, suggesting a possible role for PAI-RBP1 in tumorigenesis and tumor metastasis.

For highly specific detection of SERBP1-associated RNA regulatory signaling pathways, see our SERBP1 Antibody / mRNA Binding Protein Antibody page featuring clone SERBP1/3491 with WB, IHC, and HuProt(TM) microarray specificity validation data.

Application Notes

Optimal dilution of the SERBP1 antibody should be determined by the researcher.

1. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

Immunogen

A recombinant human partial protein corresponding to amino acids 3-139 was used as the immunogen for the SERBP1 antibody.

Storage

Store the SERBP1 antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).

Copyright © NSJ Bioreagents. All rights reserved.