


SERPINB5 Antibody / MASPIN [clone SERPINB5/4972] (V9413)


Catalog No.	Formulation	Size
V9413-100UG	0.2~mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V9413-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V9413SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	SERPINB5/4972
Purity	Protein A/G affinity
UniProt	P36952
Localization	Secreted, Extracellular space
Applications	Western Blot : 1-2ug/ml Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This SERPINB5 antibody is available for research use only.




IHC staining of FFPE human skin with SERPINB5 antibody (clone SERPINB5/4972). Negative control inset: PBS used instead of primary antibody to control for secondary Ab binding. HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human skin tissue with SERPINB5 antibody (clone SERPINB5/4972). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



Western blot testing of human PC3 cell lysate using SERPINB5 antibody (clone SERPINB5/4972). Predicted molecular weight: 42~45 kDa.



Western blot testing of human SW732 cell lysate using SERPINB5 antibody (clone SERPINB5/4972). Predicted molecular weight: 42~45 kDa.



Analysis of HuProt(TM) microarray containing more than 19,000 full-length human proteins using SERPINB5 antibody (clone SERPINB5/4972). These results demonstrate the foremost specificity of the SERPINB5/4972 mAb. Z- and S- score: The Z-score represents the strength of a signal that an antibody (in combination with a fluorescently-tagged anti-IgG secondary Ab) produces when binding to a particular protein on the HuProt(TM) array. Z-scores are described in units of standard deviations (SD's) above the mean value of all signals generated on that array. If the targets on the HuProt(TM) are arranged in descending order of the Z-score, the S-score is the difference (also in units of SD's) between the Z-scores. The S-score therefore represents the relative target specificity of an Ab to its intended target.

Description

Maspin/SERPINB5 is structurally a serine protease inhibitor (serpin) that was initially isolated from normal human mammary epithelial cells. Serpins are a family of proteins that inhibit Chymotrypsin-like serine proteinases. Serpins control activated proteinases and several are involved in the regulation of cell death. Maspin is found in the extracellular matrix and at the plasma membrane. Maspin has been shown to act at the cell surface to block cell motility and inhibit invasion of breast and prostate cancer cells. Maspin is present in normal mammary epithelial cells but is absent in many tumor cell lines, yet no major structural alterations of the Maspin gene have been identified in tumor cells. Similarly, Maspin is expressed in normal prostate cells and downregulated or absent in prostate tumor cells.

Application Notes

Optimal dilution of the SERPINB5 antibody should be determined by the researcher.

Immunogen

A portion of amino acids1-200 was used as the immunogen for the SERPINB5 antibody.

Storage Aliquot the SERPINB5 antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.