


# Sex Hormone Binding Globulin Antibody / SHBG [clone SHBG/8924] (V5419)


Catalog No.	Formulation	Size
V5419-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	100 ug
V5419-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced), 0.05% sodium azide	20 ug
V5419SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug

# **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG2b, kappa
Clone Name	SHBG/8924
Purity	Protein A/G affinity
UniProt	P04278
Localization	Cytoplasm, Extracellular (Secreted)
Applications	Immunohistochemistry (FFPE) : 1-2ug/ml
Limitations	This Sex Hormone Binding Globulin antibody is available for research use only.



IHC staining of FFPE human testis tissue with Sex Hormone Binding Globulin antibody (clone SHBG/8924). HIER: boil tissue sections in pH 9 10mM Tris with 1mM EDTA for 20 min and allow to cool before testing.



SDS-PAGE analysis of purified, BSA-free Sex Hormone Binding Globulin antibody (clone SHBG/8924) as confirmation of integrity and purity.

## **Description**

Recognizes a protein of 45kDa, identified as SHBG. It functions as an androgen transport protein, but may also be involved in receptor-mediated processes. Each dimer binds one molecule of steroid. It is specific for 5-alpha-dihydrotestosterone, testosterone, and 17-beta-estradiol. SHBG regulates the plasma metabolic clearance rate of steroid hormones by controlling their plasma concentration. In testis, it is synthesized by the Sertoli cells, secreted into the lumen of the seminiferous tubule and transported to the epididymis.

### **Application Notes**

Optimal dilution of the Sex Hormone Binding Globulin antibody should be determined by the researcher.

#### **Immunogen**

Recombinant full-length human Sex Hormone Binding Globulin protein was used as the immunogen for the Sex Hormone Binding Globulin antibody.

#### **Storage**

Aliquot the Sex Hormone Binding Globulin antibody and store frozen at -20oC or colder. Avoid repeated freeze-thaw cycles.