


# SHBG Antibody [clone SPM605] (V2854)

Catalog No.	Formulation	Size
V2854-100UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	100 ug
V2854-20UG	0.2 mg/ml in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide	20 ug
V2854SAF-100UG	1 mg/ml in 1X PBS; BSA free, sodium azide free	100 ug
V2854IHC-7ML	Prediluted in 1X PBS with 0.1 mg/ml BSA (US sourced) and 0.05% sodium azide; *For IHC use only*	7 ml

# **Bulk quote request**

Availability	1-3 business days
Species Reactivity	Human
Format	Purified
Clonality	Monoclonal (mouse origin)
Isotype	Mouse IgG1, kappa
Clone Name	SPM605
Purity	Protein G affinity chromatography
UniProt	P04278
Localization	Cytoplasmic & extracellular (secreted)
Applications	Immunohistochemistry (FFPE): 1-2ug/ml for 30 min at RT
Limitations	This SHBG antibody is available for research use only.



IHC: Formalin-fixed, paraffin-embedded human testicular carcinoma stained with SHBG antibody (clone SPM605).

## **Description**

Recognizes a protein of 45kDa, identified as SHBG. It functions as an androgen transport protein, but may also be involved in receptor-mediated processes. Each dimer binds one molecule of steroid. It is specific for 5-alpha-dihydrotestosterone, testosterone, and 17-beta-estradiol. SHBG regulates the plasma metabolic clearance rate of steroid hormones by controlling their plasma concentration. In testis, it is synthesized by the Sertoli cells, secreted into the lumen of the seminiferous tubule and transported to the epididymis.

# **Application Notes**

Optimal dilution of the SHBG antibody should be determined by the researcher.

- 1. Staining of formalin-fixed tissues requires boiling tissue sections in 10mM Tris buffer with 1mM EDTA, pH 9, for 10-20 min followed by cooling at RT for 20 min.
- 2. The prediluted format is supplied in a dropper bottle and is optimized for use in IHC. After epitope retrieval step (if required), drip mAb solution onto the tissue section and incubate at RT for 30 min.

### **Immunogen**

Recombinant full-length human protein was used as the immunogen for the SHBG antibody.

### **Storage**

Store the SHBG antibody at 2-8oC (with azide) or aliquot and store at -20oC or colder (without azide).