
Sideroflexin-3 Antibody / SFXN3 (RQ7637)


Catalog No.	Formulation	Size
RQ7637	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Antigen affinity purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	Q9BWM7
Localization	Cytoplasmic
Applications	Western Blot : 0.5-1ug/ml Immunohistochemistry (FFPE) : 2-5ug/ml Flow Cytometry : 1-3ug/million cells Immunofluorescence : 5ug/ml Direct ELISA : 0.1-0.5ug/ml
Limitations	This Sideroflexin-3 antibody is available for research use only.




Immunofluorescent staining of FFPE human renal cancer tissue with Sideroflexin-3 antibody (green) and DAPI nuclear stain (blue). HIER: steam section in pH8 EDTA buffer for 20 min.




IHC staining of FFPE human spleen tissue with Sideroflexin-3 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human breast cancer tissue with Sideroflexin-3 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human liver cancer tissue with Sideroflexin-3 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human thyroid cancer tissue with Sideroflexin-3 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human liver cancer tissue with Sideroflexin-3 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



IHC staining of FFPE human lung adenocarcinoma tissue with Sideroflexin-3 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Western blot testing of human 1) HeLa and 2) U-87 MG cell lysate with Sideroflexin-3 antibody. Predicted molecular weight ~36 kDa.



Flow cytometry testing of human A549 cells with Sideroflexin-3 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= Sideroflexin-3 antibody.

Description

SFXN3, a mitochondrial membrane protein, is a member of sideroflexin family. Serum anti-SFXN3 autoantibody was elevated in early stage of the oral squamous cell carcinoma significantly, indicating its clinical value for diagnosis of oral squamous cell carcinoma.

Application Notes

Optimal dilution of the Sideroflexin-3 antibody should be determined by the researcher.

Immunogen

E. coli-derived recombinant human protein (amino acids D32-L321) was used as the immunogen for the Sideroflexin-3 antibody.

Storage

After reconstitution, the Sideroflexin-3 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.

Ordering: Phone: 858.663.9055 | Fax: 1.267.821.0800 | Email: info@nsjbio.com
Copyright © NSJ Bioreagents. All rights reserved.