
SITAC Antibody / SDCBP2 / Syntenin-2 (RQ7612)


Catalog No.	Formulation	Size
RQ7612	0.5mg/ml if reconstituted with 0.2ml sterile DI water	100 ug

Bulk quote request


Availability	1-3 business days
Species Reactivity	Human
Format	Antigen affinity purified
Host	Rabbit
Clonality	Polyclonal (rabbit origin)
Isotype	Rabbit IgG
Purity	Antigen affinity purified
Buffer	Lyophilized from 1X PBS with 2% Trehalose
UniProt	Q9H190
Localization	Cytoplasmic, nuclear
Applications	Western blot: 0.5-1ug/ml Immunohistochemistry (FFPE): 2-5ug/ml Flow cytometry: 1-3ug/million cells Immunofluorescence: 5ug/ml Direct ELISA: 0.1-0.5ug/ml
Limitations	This SDCBP2 antibody is available for research use only.




Immunofluorescent staining of FFPE human U-2 OS cells with SDCBP2 antibody (green) and Beta Tubulin mAb (red). HIER: steam section in pH6 citrate buffer for 20 min.




IHC staining of FFPE human colon adenocarcinoma tissue with SDCBP2 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human liver cancer tissue with SDCBP2 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human breast cancer tissue with SDCBP2 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human renal cancer tissue with SDCBP2 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human lung adenocarcinoma tissue with SDCBP2 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human prostate adenocarcinoma tissue with SDCBP2 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.




IHC staining of FFPE human spleen tissue with SDCBP2 antibody. HIER: boil tissue sections in pH8 EDTA for 20 min and allow to cool before testing.



Western blot testing of human 1) A431 and 2) HaCaT cell lysate with SDCBP2 antibody. Predicted molecular weight ~32 kDa, commonly observed at 32-37 kDa with a possible ~23 kDa isoform.



Flow cytometry testing of human MCF7 cells with SDCBP2 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= SDCBP2 antibody.



Flow cytometry testing of human U-2 OS cells with SDCBP2 antibody at 1ug/million cells (blocked with goat sera); Red=cells alone, Green=isotype control, Blue= SDCBP2 antibody.

Description

Syntenin-2 is a protein that in humans is encoded by the SDCBP2 gene. The protein encoded by this gene contains two class II PDZ domains. PDZ domains facilitate protein-protein interactions by binding to the cytoplasmic C-terminus of transmembrane proteins, and PDZ-containing proteins mediate cell signaling and the organization of protein complexes. The encoded protein binds to phosphatidylinositol 4, 5-bisphosphate (PIP2) and plays a role in nuclear PIP2 organization and cell division. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene. Read-through transcription also exists between this gene and the upstream FKBP1A (FK506 binding protein 1A, 12kDa) gene, as represented in GeneID:100528031.

Application Notes

Optimal dilution of the SDCBP2 antibody should be determined by the researcher.

Immunogen

E. coli-derived recombinant human protein (amino acids D14-A292) was used as the immunogen for the SDCBP2 antibody.

Storage

After reconstitution, the SDCBP2 antibody can be stored for up to one month at 4oC. For long-term, aliquot and store at -20oC. Avoid repeated freezing and thawing.

Copyright © NSJ Bioreagents. All rights reserved.